

IZA DP No. 7657

The Origins of Early Childhood Anthropometric Persistence

Daniel L. Millimet
Rusty Tchernis

September 2013

The Origins of Early Childhood Anthropometric Persistence

Daniel L. Millimet
*Southern Methodist University
and IZA*

Rusty Tchernis
*Georgia State University,
NBER and IZA*

Discussion Paper No. 7657
September 2013

IZA

P.O. Box 7240
53072 Bonn
Germany

Phone: +49-228-3894-0
Fax: +49-228-3894-180
E-mail: iza@iza.org

Any opinions expressed here are those of the author(s) and not those of IZA. Research published in this series may include views on policy, but the institute itself takes no institutional policy positions. The IZA research network is committed to the IZA Guiding Principles of Research Integrity.

The Institute for the Study of Labor (IZA) in Bonn is a local and virtual international research center and a place of communication between science, politics and business. IZA is an independent nonprofit organization supported by Deutsche Post Foundation. The center is associated with the University of Bonn and offers a stimulating research environment through its international network, workshops and conferences, data service, project support, research visits and doctoral program. IZA engages in (i) original and internationally competitive research in all fields of labor economics, (ii) development of policy concepts, and (iii) dissemination of research results and concepts to the interested public.

IZA Discussion Papers often represent preliminary work and are circulated to encourage discussion. Citation of such a paper should account for its provisional character. A revised version may be available directly from the author.

ABSTRACT

The Origins of Early Childhood Anthropometric Persistence¹

Rates of childhood obesity have increased dramatically in the last few decades. Non-causal evidence suggests that childhood obesity is highly persistent over the life cycle. However little is known about the origins of this persistence. In this paper we attempt to answer three questions. First, how do anthropometric measures evolve from birth through primary school? Second, what is the *causal* effect of past anthropometric outcomes on future anthropometric outcomes? In other words, how important is *state dependence* in the evolution of anthropometric measures during the early part of the life cycle. Third, how important are time-varying and time invariant factors in the dynamics of childhood anthropometric measures? We find that anthropometric measures are *highly persistent* from infancy through primary school. Moreover, most of this persistence is driven by unobserved, time invariant factors that are determined prior to birth, consistent with the so-called fetal origins hypothesis. As such, interventions designed to improve child anthropometric status will only have meaningful, long-run effects if these time invariant factors are altered. Unfortunately, future research is needed to identify such factors, although evidence suggests that maternal nutrition may play an important role.

JEL Classification: C23, I12, I18

Keywords: childhood obesity, persistence, fetal origins hypothesis

Corresponding author:

Daniel L. Millimet
Department of Economics
Southern Methodist University
Box 0496
Dallas, TX 75275-0496
USA
E-mail: millimet@smu.edu

¹ This study was conducted by Georgia State University and Southern Methodist University under a cooperative agreement with the U.S. Department of Agriculture, Economic Research Service, Food and Nutrition Assistance Research Program (agreement no. 58-5000-0-0080). The views expressed here are those of the authors and do not necessarily reflect those of the USDA or ERS. The authors are grateful to Chris Ruhm for helpful comments.

1 Introduction

The rise in childhood obesity in the U.S. is well chronicled. Moreover, strong evidence suggests that childhood obesity is highly persistent over the life cycle. However little is known about the origins of this persistence. In this paper we attempt to answer three questions. First, how do weight and height evolve from birth through primary school? Second, what is the *causal* effect of past weight and height status on the future weight and height of children? In other words, how important is *state dependence* in the evolution of anthropometric measures during the early part of the life cycle. Third, how important are time-varying and time invariant factors in the dynamics of childhood weight and height?

These are important public health questions as the prevalence of obese adolescents has tripled in the last thirty years; it has more than doubled for younger children. Defined as having an age- and sex-adjusted body mass index (BMI) above the 95th percentile of the reference distribution, the prevalence of obese children increased from 5% to 12.4% for 2-5 year old children and from 5% to 17.6% for 12 to 19 year-olds between 1976 and 2006 (Ogden et al. 2008). In addition, vast differences in the time trends of BMI increases have been documented: the incidence of obesity among white girls aged 12-19 has increased from 7.4% to 14.5% between 1988 and 2006, whereas the corresponding figures for African-American girls are 13.2% and 27.7% (Ogden et al. 2002; Ogden et al. 2008). Deckelbaum and Williams (2001, p. 242S) conclude that “childhood obesity is increasing at epidemic rates, even among pre-school children...” More recently, Brisbois et al. (2012, p. 347) state: “Obesity is considered to be a worldwide epidemic with little evidence that its incidence is declining or that it has even reached a plateau.”

While there exists some evidence that childhood obesity rates may have begun to recede in the U.S., concern over childhood obesity remains high due to the well-documented consequences of obesity and the lack of understanding pertaining to the turnaround.² Obesity burdens individuals with severe physical, economic, and emotional suffering, and puts children and adolescents at risk for a number of health problems such as those affecting cardiovascular health, the endocrine system, and mental health (Deckelbaum and Williams 2001; Krebs and Jacobson 2003). Dietz and Gortmaker (2001) note that 60% of overweight children aged five to ten years old have at least one associated cardiovascular disease risk factor.

Perhaps the largest cost of childhood obesity comes from its impact on adult obesity. Currently, 60% of the total U.S. population is overweight or obese and 50% is expected to be obese in 2030 at the current rate (Dor et al. 2010). Walpole et al. (2012) calculate that North America accounts for 34% of the total human biomass in the world despite containing only 6% of the world population. Moreover, the authors estimate that if the entire world had the same BMI distribution as the U.S., this would be equivalent to an additional 935 million people in the world of average BMI. Finkelstein and Zuckerman (2007) report that if the childhood obesity epidemic continues unabated at the current rate, as many as 30-40% of the US population will develop Type 2 Diabetes during their lifetime. Mocan and Tekin (2011) document the links between adult obesity and lower wages, productivity, and self-esteem. In the U.S., the total cost attributable to obesity was over \$75 billion in 2000 according to Finkelstein et al. (2004). More recent estimates put the cost over \$200 billion (Cawley and Meyerhoefer 2012).

While the changes in childhood obesity rates across cohorts, as well as the consequences of these increases, are

²See <http://www.nytimes.com/2013/08/07/health/broad-decline-in-obesity-rate-seen-in-poor-young-children.html?pagewanted=all>.

well-documented, much less is known about how anthropometric measures of children evolve over the life cycle. A growing literature has investigated persistence in anthropometric outcomes in a non-causal framework, stressing the correlation in outcomes over time.³ Whitaker et al. (1997) found that the probability of an overweight six year-old child becoming an obese adult is 50% compared to 10% for a non-overweight child. In addition, the risk of becoming obese in adulthood is exacerbated by having an obese parent. Eriksson et al. (2001) found that individuals were three times more likely to be obese as an adult if they had a BMI greater than 16, as opposed to below 14.5, at age seven. Nader et al. (2007) find that children who were overweight prior to the age of five are five times as likely to be overweight at 12 relative to children who were not overweight prior to the age of five.

Freedman et al. (2001) also report a strong relationship between overweight status in childhood and adult BMI. However, most striking is that obese adults who were overweight prior to age eight have a much higher BMI than individuals suffering from adult onset obesity (41 versus 35). In a later study, Freedman et al. (2005) document significant differences in the transmission of BMI from childhood to adulthood along racial lines. Gable et al. (2008) analyze the relationship between socioeconomic status, overweight persistence, and school outcomes. The authors find that family socioeconomic status is predictive of both the probability of a child being overweight and the probability of persistence of overweight status. Van Cleave et al. (2010) analyze changes in the prevalence of obesity and other chronic conditions (e.g., asthma, other physical and learning conditions). The authors find that prevalence of obesity is increasing and is highly persistent over time. Conversely, many children with chronic conditions at ages two through eight did not have the condition six years later. Finally, Millimet and Tchernis (2013) assess persistence during infancy and primary school, documenting a significant increase in persistence upon entry into primary school. Deckelbaum and Williams (2001, p. 239S) conclude: “Disturbingly, obesity in childhood, particularly in adolescence is a key predictor for obesity in adulthood.” Similarly, Dietz and Gortmaker (2001, p. 340) state: “The best evidence suggests that the majority of overweight adolescents go on to be overweight adults.”

We build on this prior literature in an attempt to uncover the origins of the persistence in anthropometric measures. Specifically, we revisit the question of persistence in early childhood health outcomes and investigate the relative importance of state dependence (i.e., a causal effect of past anthropometric status on future status), unobserved heterogeneity (i.e., unobserved genetic or environmental risk factors), and observed heterogeneity (i.e., commonly measured risk factors) on this persistence. We then ask whether the origins of anthropometric persistence vary by age, race, gender, or socioeconomic status. Our analysis is fundamentally important for researchers as well as policymakers. If obesity has its origins early in life and is persistent over time, then early intervention is preferable to waiting until adolescence or beyond.⁴ However, and perhaps most importantly, if persistence is due to persistent underlying factors rather than state dependence, then only by altering these factors can children be moved to a different trajectory.

³Iughetti et al. (2008) provide an excellent review.

⁴For instance, an article in the *New York Times* on March 22, 2010 states that some evidence now suggests that children may become entrenched “on an obesity trajectory” even before kindergarten; however, the evidence is not “ironclad” (<http://www.nytimes.com/2010/03/23/health/23obese.html>). Public health officials tend to advocate school-based reforms in light of the near universal enrollment, yet others stress the importance of preschool interventions (e.g., Frisvold and Giri 2011; Dietz and Gortmaker 2001; Davis and Christoffel 1994). Eriksson et al. (2001, p. 735) conclude that “obesity is initiated early in life.”

To examine these fundamental questions, we estimate dynamic regression models using data from the Early Childhood Longitudinal Survey – Kindergarten Cohort (ECLS-K). The ECLS-K is a nationally representative longitudinal survey of children entering kindergarten in Fall 1998. In addition to providing information on birthweight, anthropometric data is collected at several points in time between kindergarten and eighth grade. We then supplement this analysis by examining data from the Early Childhood Longitudinal Survey – Birth Cohort (ECLS-B). The ECLS-B is a nationally representative longitudinal survey of children born in the U.S. in 2001. Information is provided on these children at ages 9 months, two years, four years, and five years. Thus, the ECLS-B sample allows for a more refined examination of anthropometric trajectories prior to kindergarten entry.

The analysis leads to two salient conclusions. First, weight, height, and BMI are *highly persistent* starting in *early* infancy. Second, the vast majority of persistence is attributable to *time invariant* characteristics of children. This finding is of critical importance as it implies that the only interventions that will have a substantive, long-run effect on a child’s anthropometric status are those that alter these salient, time invariant attributes. Thus, current policy interventions may, at best, have a marginal impact in the short-run and, at worst, be destined to fail (see, e.g., Davis and Gebremariam (2010)). Moreover, while it is difficult to say what these critical, time invariant attributes are given the data at hand, we find some evidence that fetal nutrition – as proxied by mother’s pre-pregnancy weight and weight gain during pregnancy, gestation age, birth status (singleton, twin, or higher order birth), and birthweight – impacts the evolution of anthropometric measures over the early life cycle. However, *unobserved*, time invariant attributes play a much more prominent role.

The notion that attributes determined at or shortly after birth, and thus time invariant over the life of an individual, play a dominant role in the evolution of obesity is *consistent* with the strong evidence in economics and elsewhere on the so-called fetal origins hypothesis (see, e.g., Almond and Currie 2011).⁵ The fetal origins hypothesis, also referred to as the thrifty phenotype hypothesis or Barker’s hypothesis (due to Barker’s original publication in 1992), posits long-run effects of conditions *in utero* during critical periods of development through “programmed” changes in the physiology and metabolism of individuals (Barker 1997). An article in *Time* on September 22, 2010 summarizes⁶:

“[P]ioneers assert that the nine months of gestation constitute the most consequential period of our lives, permanently influencing the wiring of the brain and the functioning of organs such as the heart, liver and pancreas. The conditions we encounter in utero, they claim, shape our susceptibility to disease, our appetite and metabolism, our intelligence and temperament. In the literature on the subject, which has exploded over the past 10 years, you can find references to the fetal origins of cancer, cardiovascular disease, allergies, asthma, hypertension, diabetes, obesity, mental illness — even of conditions associated with old age like arthritis, osteoporosis and cognitive decline.”

Our analysis is consistent with this view, the implications of which are quite profound. If correct, the most efficient interventions to curb obesity may need to start *prior* to childbirth. Deckelbaum & Williams (2001, p. 239S) conclude:

⁵While the findings here are consistent with the fetal origins hypothesis, we cannot eliminate other possible explanations for what these salient, unobserved attributes entail.

⁶See <http://www.time.com/time/magazine/article/0,9171,2021065,00.html>.

“Novel approaches in the prevention and treatment of childhood overweight and obesity are urgently required. With the strong evidence that a lifecycle perspective is important in obesity development and its consequences, consideration must be focused on prevention of obesity in women of child-bearing age, excessive weight gain during pregnancy, and the role of breast-feeding in reducing later obesity in children and adults. Consideration must be given to family behavior patterns, diet after weaning, and the use of new methods of information dissemination to help reduce the impact of childhood obesity worldwide.”

The remainder of the paper is organized as follows. Section 2 provides a brief overview of the prior literature. Section 3 presents the empirical methodology and data overview. Section 4 discusses the results and their implications. Section 5 concludes.

2 Related Literature

In addition to the literature already discussed pertaining to the correlations between childhood weight status and adult obesity, two other prior strands of literature are worth discussing. The first strand includes investigations on the persistence in health among adolescents and adults in a *causal* framework. For example, Halliday (2008) investigates persistence in self-reported health status among white adults age 22-60 using data from the PSID and allows the parameters of the model to vary. The results suggest that the degree of state dependence – the causal effect of past states on one’s current state – in health is modest for half the population, yet it explains much of the observed persistence in health for the other half. Ham et al. (2013) analyze persistence in bulimia nervosa in young women. The authors find a substantial role for state dependence in the persistence of bulimia nervosa, thus justifying the importance of early intervention. Our analysis follows the logic of these studies.

The second strand focuses explicitly on the fetal origins hypothesis. As stated earlier, beginning with Barker’s work, there is a strong belief that *in utero* events may determine whether a fetus ends up on an “obesity trajectory.” Deckelbaum & Williams (2001, p. 239S) note that “emerging data suggest associations between the influence of maternal and fetal factors during intrauterine growth and growth during the first year of life, on risk of later development of adult obesity and its comorbidities.” More recently, Brisbois et al. (2012, p. 347) state: “Based on recent evidence, early-life experiences *in utero* and postnatal influences may induce permanent changes in physiologic function that programme the long-term regulation of energy balance. This subsequently may adversely impact obesity risk in later life.”

Which factors may induce such permanent changes in order to set a fetus upon an “obesity trajectory” is the subject of on-going research. While initial hypotheses focused on undernutrition and oxygen supply, additional factors such as maternal BMI, maternal weight gain, maternal smoking, gestational diabetes requiring insulin, and postnatal characteristics such as breastfeeding and the timing of introduction to solid foods are also found to be important (Dietz 1997; Deckelbaum and Williams 2001; Brisbois et al. 2012).

Within this literature, studies have also focused on the identification of early life physical indicators of predisposition to future obesity. Preliminary results suggest that birthweight, length, and gestation age at birth alone are not strong predictors. Instead, there are complex interactions between these measures, along with other measures such

as head circumference, that matter. For example, a fetus born prematurely and, as a result, with low birthweight and length is not likely to be at greater risk of future obesity as long as the fetus’ measurements are in proportion and within ‘normal’ ranges given its gestation age. On the other hand, a fetus born with disproportionate physical measurements suggests a greater risk (Barker 1997; Sayer et al. 1997; Godfrey and Barker 2001; Brisbois et al. 2012).

3 Empirics

3.1 Methodology

We assess the extent and origins of persistence using a dynamic regression framework. This approach allows for the decomposition of persistence into various components reflecting state dependence, observed heterogeneity, and unobserved heterogeneity.

The simplest estimating equation is

$$y_{it} = \gamma y_{it-1} + \varepsilon_{it}, \quad i = 1, \dots, N; t = 1, \dots, T \tag{1}$$

where y_{it} is a measure of weight status for child i at time t , ε_{it} is a mean zero error term, and T must be at least two (given observability of the initial observation, y_{i0}). Here, γ reflects the overall level of persistence as it captures the entire association between past and current anthropometric status. To decompose this overall persistence, we next incorporate *observed* heterogeneity into the model

$$y_{it} = \gamma y_{it-1} + x_{it}\beta + w_i\delta + \varepsilon_{it}, \quad i = 1, \dots, N; t = 1, \dots, T \tag{2}$$

where x_{it} is a vector of observed, time-varying attributes of child i at time t and w_i is a vector of observed, time invariant attributes of child i . The change in the estimate of γ from (1) to (2) reflects the portion of persistence attributable to persistent, observed heterogeneity. Finally, we include observed time-varying heterogeneity and all sources (observed and unobserved) of time-invariant heterogeneity into the model

$$y_{it} = \gamma y_{it-1} + x_{it}\beta + \alpha_i + \varepsilon_{it}, \quad i = 1, \dots, N; t = 1, \dots, T \tag{3}$$

where α_i is a child-specific fixed effect. In (3) γ reflects the degree of *state dependence* as it captures the causal effect of past weight status on current weight status. The child-specific fixed effect, α , reflects persistence in child anthropometric measures due to persistent observed and unobserved, child-specific heterogeneity (such as time invariant environmental and genetic factors). In such models, β represents the contemporaneous effects of the observed, time varying regressors, whereas $\beta/(1 - \gamma)$ represents the long-run effects of a permanent unit change in these variables.

Estimation of (3) is straightforward (assuming the model is correctly specified). Following Anderson and Hsiao (1981), (3) is first-differenced to eliminate α_i . The first-differenced model is then estimated via instrumental variables since the first-differenced lagged dependent variable is necessarily correlated with the first-differenced error term. However, y_{it-2} represents a valid instrument if ε is serially uncorrelated. The models are estimated by Generalized Method of Moments (GMM).

Once the models are estimated, in addition to simply examining the coefficient estimates, we follow the logic in Ulrick (2008) and simulate probabilities such as the following

$$\Pr(y_{iT} \geq y^* | y_{i0} \geq y_0) \quad (4)$$

given estimates of the regression model. Here, (4) represents the probability of a child having an anthropometric measure above y^* in the terminal period conditional on an initial measure greater than or equal to some value y_0 . For example, one might be interested in the probability of a child having a BMI above the 85th percentile in period T conditional on being above the 85th percentile in the initial period, $t = 0$. These probabilities incorporate not just the coefficient directly related to persistence, γ , but also reflect persistence due to persistence in observed and unobserved determinants of child weight. Moreover, by altering the attributes of individuals, we can simulate counterfactual probabilities as well. Finally, we can simulate these probabilities and counterfactual probabilities for different socioeconomic groups. This allows one to determine if the degree of persistence, and the factors contributing to such persistence, vary across socioeconomic groups.

Before detailing the simulations undertaken, note that upon estimating (3), estimates of α_i are given by

$$\hat{\alpha}_i = \frac{1}{T} \sum_{t=1}^T \left[y_{it} - \hat{\gamma} y_{it-1} - x_{it} \hat{\beta} \right], \quad i = 1, \dots, N. \quad (5)$$

The estimates can then be decomposed into observed and unobserved time invariant factors by estimating the following model using OLS

$$\hat{\alpha}_i = w_i \delta + \eta_i, \quad (6)$$

where w_i now includes an intercept and η is a mean zero error term. Finally, given estimates of γ , β , and α , we can obtain estimates of the idiosyncratic errors, ε and η , using (3) and (6).

To proceed, we simulate probabilities, such as those given by (4), under the following counterfactual scenarios:

1. Own y_{i0} , own x_{it} , set $\varepsilon_{it} = 0$, and
 - (a) replace $\alpha_i = \bar{\alpha}$, or
 - (b) draw $\alpha_i \sim F(\alpha)$ where $F(\cdot)$ is the empirical distribution of α_i , or
 - (c) draw $\alpha_i \sim F_G(\alpha)$ where $F_G(\cdot)$ is the empirical distribution of α_i in sub-sample G and $i \in G$, or
 - (d) draw $\alpha_i \sim F_{G'}(\alpha)$ where $F_{G'}(\cdot)$ is the empirical distribution of α_i in sub-sample G' and $i \notin G'$.
2. Own y_{i0} , own x_{it} , set $\eta_i = 0$, set $\varepsilon_{it} = 0$, and
 - (a) own w_i , or
 - (b) draw $w_i \sim F(w)$ where $F(\cdot)$ is the empirical distribution of w_i , or
 - (c) draw $w_i \sim F_G(w)$ where $F_G(\cdot)$ is the empirical distribution of w_i in sub-sample G and $i \in G$, or
 - (d) $w_i \sim F_{G'}(w)$ where $F_{G'}(\cdot)$ is the empirical distribution of w_i in sub-sample G' and $i \notin G'$.
3. Own y_{i0} , own x_{it} , own w_i , set $\varepsilon_{it} = 0$, and

- (a) draw $\eta_i \sim F(\eta)$ where $F(\cdot)$ is the empirical distribution of η_i , or
 - (b) draw $\eta_i \sim F_G(\eta)$ where $F_G(\cdot)$ is the empirical distribution of η_i in sub-sample G and $i \in G$, or
 - (c) draw $\eta_i \sim F_{G'}(\eta)$ where $F_{G'}(\cdot)$ is the empirical distribution of η_i in sub-sample G' and $i \notin G$.
4. Own y_{i0} , own α_i , set $\varepsilon_{it} = 0$, and
- (a) replace $x_{it} = \bar{x}_t$, or
 - (b) draw $x_i \sim F(x_1, \dots, x_T)$ where $F(\cdot)$ is the empirical joint distribution of x_1, \dots, x_T , or
 - (c) draw $x_i \sim F_G(x_1, \dots, x_T)$ where $F_G(\cdot)$ is the empirical joint distribution of x_1, \dots, x_T in sub-sample G and $i \in G$, or.
 - (d) draw $x_i \sim F_{G'}(x_1, \dots, x_T)$ where $F_{G'}(\cdot)$ is the empirical joint distribution of x_1, \dots, x_T in sub-sample G and $i \in G$.
5. Own y_{i0} , own x_{it} , own α_i , and
- (a) draw $\varepsilon_i \sim F(\varepsilon_1, \dots, \varepsilon_T)$ where $F(\cdot)$ is the empirical distribution of ε_i , or
 - (b) draw $\varepsilon_i \sim F_G(\varepsilon_1, \dots, \varepsilon_T)$ where $F_G(\cdot)$ is the empirical distribution of ε_i in sub-sample G and $i \in G$, or
 - (c) draw $\varepsilon_i \sim F_{G'}(\varepsilon_1, \dots, \varepsilon_T)$ where $F_{G'}(\cdot)$ is the empirical distribution of ε_i in sub-sample G' and $i \notin G$.
6. Own y_{i0} , own α_i , and
- (a) draw $x_i, \varepsilon_i \sim F(x_1, \dots, x_T, \varepsilon_1, \dots, \varepsilon_T)$ where $F(\cdot)$ is the empirical joint distribution of $x_1, \dots, x_T, \varepsilon_1, \dots, \varepsilon_T$, or
 - (b) draw $x_i, \varepsilon_i \sim F_G(x_1, \dots, x_T, \varepsilon_1, \dots, \varepsilon_T)$ where $F_G(\cdot)$ is the empirical joint distribution of $x_1, \dots, x_T, \varepsilon_1, \dots, \varepsilon_T$ in sub-sample G and $i \in G$, or
 - (c) draw $x_i, \varepsilon_i \sim F_{G'}(x_1, \dots, x_T, \varepsilon_1, \dots, \varepsilon_T)$ where $F_{G'}(\cdot)$ is the empirical joint distribution of $x_1, \dots, x_T, \varepsilon_1, \dots, \varepsilon_T$ in sub-sample G' and $i \notin G$.

Probabilities are obtained using 500 simulations. See the Appendix A for further details.

Case 1 eliminates time-varying, unobserved heterogeneity, ε_{it} , and assesses the impact of altering the distribution of time invariant heterogeneity, α_i . Case 1a eliminates all time invariant heterogeneity. Cases 1b-1d replace actual time invariant heterogeneity with a random draw. Case 1b draws from the empirical distribution. Case 1c draws from the empirical distribution of the same sub-group as observation i . Case 1d draws from the empirical distribution of the sub-group to which observation i does not belong. For example, if we divide the sample based on gender, Case 1c draws a value of α from the empirical distribution of boys for each boy. Case 1d entails drawing a value of α from the empirical distribution of girls for each boy. Case 1b succeeds in entirely breaking any correlation between the initial condition, y_{i0} , and x_{it} and time invariant heterogeneity, α_i . Case 1c partially breaks this correlation. In total, these cases speak to the relative importance of time invariant heterogeneity in the persistence of weight status, as well as differences in the distribution of α across different sub-groups.

Case 2 continues to eliminate time-varying, unobserved heterogeneity, ε_{it} . However, time invariant, unobserved heterogeneity, η_i , is now also eliminated; the observed component of time invariant heterogeneity is then altered. Case 2a utilizes each observation’s own time invariant heterogeneity, w_i . Case 2b draws w_i from the empirical distribution. Case 2c draws w_i from the empirical distribution of the same sub-group as observation i . Case 2d draws w_i from the empirical distribution of the sub-group to which observation i does not belong. Case 3 is similar, but has individuals retain their time invariant, observed heterogeneity, w_i , and alters the distribution of time invariant, unobserved heterogeneity, η_i . Case 3a draws η_i from the population empirical distribution. Case 3b draws η_i from the empirical distribution of the same sub-group as observation i . Case 3c draws η_i from the empirical distribution of the sub-group to which observation i does not belong. Altogether, Cases 2 and 3 permit assessment of the relative importance of the observed and unobserved components of time invariant heterogeneity in the persistence of weight status. Moreover, they will also illuminate any salient differences in these components across different sub-groups.

Case 4 continues to eliminate time-varying, unobserved heterogeneity, ε_{it} , and assesses the impact of altering the distribution of time-varying, observed heterogeneity, x_{it} . Case 4a eliminates all time-varying heterogeneity. Cases 4b-4d replace actual time-varying, observed heterogeneity with a random draw. Case 4b draws from the empirical distribution. Case 4c draws from the empirical distribution of the same sub-group as observation i . Case 4d draws from the empirical distribution of the sub-group to which observation i does not belong. Case 4b succeeds in entirely breaking any correlation between the initial condition, y_{i0} , and α_i and time-varying, observed heterogeneity, x_{it} . Case 4c partially breaks this correlation. These cases complement the simulations performed in Case 1 as they speak to the relative importance of time-varying, observed heterogeneity in the persistence of weight status, as well as differences in the distribution of x across different sub-groups.

Case 5 has individuals retain their time-varying, observed attributes, x_{it} , and time invariant attributes, α_i and y_{i0} , but alters the distribution of time-varying, unobserved heterogeneity, ε_{it} . Case 5a draws ε_i from the empirical distribution. Case 5b draws ε_i from the empirical distribution of the same sub-group as observation i . Case 5c draws ε_i from the empirical distribution of the sub-group to which observation i does not belong. Finally, Case 6 has individuals only retain their time invariant attributes, α_i and y_{i0} . All time-varying heterogeneity is sampled. Case 6a draws x_i and ε_i from the population empirical distribution. Case 6b draws x_i and ε_i from the empirical distribution of the same sub-group as observation i . Case 6c draws x_i and ε_i from the empirical distribution of the sub-group to which observation i does not belong. Thus, these final two cases address the relative importance of the observed and unobserved components of time-varying heterogeneity in the persistence of weight status. The results will also highlight any important differences in these components across different sub-groups.

3.2 Data

We utilize data from the restricted version of the ECLS-K. Collected by the US Department of Education, the ECLS-K surveys a nationally representative cohort of children throughout the US in fall and spring kindergarten, fall and spring first grade, spring third grade, spring fifth grade, and spring eighth grade. The sample includes data on over 20,000 students who entered kindergarten in one of roughly 1,000 schools during the 1998-99 school year. In addition to family background information, height and weight measures are available from children in each round,

as well as information on birth weight.

Our final sample consists of children for whom we have valid measures of age, gender, height, and weight.⁷ From the information on height and weight of the children, we obtain z -scores for weight, height, and BMI. Note that z -scores and percentiles are based on CDC 2000 growth charts; these are age- and gender-specific, are adjusted for normal growth, and percentiles are based on the underlying reference population.⁸ The estimation utilizes data from five waves: fall kindergarten, spring first grade, spring third grade, spring fifth grade, and spring eighth grade.⁹ The sample is a balanced a panel of roughly 9,160 children.¹⁰

Data on family background are used in two different manners in the analysis. First, we define different demographic groups in order to split the sample during the probability simulations. We consider five different partitions based on race (white vs. non-white), gender (male vs. female), urban status (urban vs. rural/suburban), mother’s education (college vs. less than college), and socioeconomic status (low vs. high SES). Second, we incorporate time-varying, x_{it} , and time invariant, w_i , attributes into the regression model.

The following time invariant covariates are included: gender, race/ethnicity (white, black, Hispanic, Asian, and other), birthweight, indicator for premature birth, indicator for being born in the U.S., indicator for being a native English speaker, city type (urban, suburban, or rural), region (northeast, midwest, south, and west), mother’s education (less than high school, high school/GED, some college, four-year college degree, and more than four years of college), mother’s age at first birth, mother’s marital status at birth, indicator for attending nonparental pre-kindergarten, indicator for mother’s labor force participation during infancy, indicator for mother’s participation in WIC (Women’s, Infants, and Children) during pregnancy, indicator for mother’s participation in WIC during infancy, indicator for mother’s participation in TANF (Temporary Assistance for Needy Families) during infancy, indicator for participation in FSP (Food Stamp Program) during infancy, and indicator for attending full day kindergarten.¹¹ The following time-varying covariates are included: an index of SES status, indicator for the household being in poverty, number of children’s books in the household, household size, family type (two parents plus siblings, two parents and no siblings, one parent and siblings, one parent and no siblings, and other), mother’s labor force status (full-time, part-time, and not working), indicator for mother absent from the household, indicator of current TANF participation, indicator of current FSP participation, indicator for health insurance, hours spent watching television during the school week, hours spent watching television during the weekend, indicator for household rules regarding television watching, days per week household eats breakfast together, days per week household eats dinner together, indicator for household food security (household never worried about running out of food), neighborhood safety (very safe, somewhat safe, and not safe), and percent of minority students in class at school. For all covariates (except

⁷The initial sample size of the ECLS-K is 21,260. After cleaning age, weight, and height as described in Millimet and Tchernis (2012, Appendix C), and due to sample attrition, the sample size falls to 9,360 in the final wave of the data. Restricting the sample to a balanced panel reduces the sample size to approximately 9,160. This is the final sample size per wave in the analysis. Note, all sample sizes are rounded to the nearest 10 per NCES restricted data regulations.

⁸ z -scores and their percentiles are obtained using the *-zanthro-* command in Stata.

⁹The survey design is troublesome in that the ECLS-K contains irregularly spaced waves. To minimize the issue, we omit the spring kindergarten wave and thus each period conceptually represents roughly a two-year window.

¹⁰Sample sizes are rounded to the nearest 10 per NCES restricted data regulations.

¹¹FSP was renamed the Supplemental Nutrition Assistance Program (SNAP) in October 2008. Since the data pre-dates this change, we refer to the program as FSP.

gender, age, height, and weight), we include dummy variables for missing observations.

4 Results

Tables 1, 5, and 9 display the results from estimation of (1), (2), and (3) for weight, height, and BMI z -scores, respectively. In addition to reporting estimates of the coefficient on the lagged outcome, γ , we report the first-stage Kleibergen-Paap (2006) Wald rk F -statistic, the Kleibergen-Paap (2006) rk test of underidentification, and a test of endogeneity. The first two tests are designed to detect any issues associated with weak instruments. Finally, recall that within each sample (i.e., the overall sample of demographic sub-group), the estimate of γ from (1) reflects the overall level of persistence, the change in the estimate moving from (1) from (2) captures the portion of persistence explained by the observable covariates, and the change moving from (2) to (3) reflects the portion of persistence explained by time invariant, observed factors.

The remaining tables present the dynamic simulations based on (4) to provide further analysis of the sources of persistence, the role of time-varying and time invariant observed attributes, and differences across demographic groups. As noted earlier, the simulations are based on the estimates of the fixed effects specification given in (3), along with the subsequent estimates of the fixed effects and their decomposition given in (5) and (6). For each outcome, we simulate three sets of probabilities:

1. $\Pr(y_{iT} \geq 85^{\text{th}} \text{ percentile} \mid y_{i0} \geq 85^{\text{th}} \text{ percentile})$,
2. $\Pr(y_{iT} \geq 95^{\text{th}} \text{ percentile} \mid y_{i0} \geq 95^{\text{th}} \text{ percentile})$, and
3. $\Pr(y_{iT} \geq 85^{\text{th}} \text{ percentile} \mid y_{i0} \leq 50^{\text{th}} \text{ percentile})$,

where period T denotes spring eighth grade and period 0 corresponds to fall kindergarten. Note, the percentile outcomes are based on the underlying reference population used in the CDC 2000 growth charts, *not* the current sample. Thus, the 85th and 95th percentiles correspond to usual cutoffs for overweight and obese when examining BMI. Finally, each table presents the *benchmark* probability, which is the empirical probability observed in the data (i.e., the sample probability as opposed to an estimate), for comparison.

4.1 Weight

Table 1 displays the regression results for weight z -scores. For the full sample, the estimates of γ across the three specifications are 0.931, 0.932, and 0.775 (standard errors are 0.003, 0.003, and 0.067, respectively). Each is statistically significant at the $p < 0.01$ confidence level and all three specifications are strongly identified. The fact that the estimate of γ does not change moving from (1) to (2) implies that our lengthy vector of time-varying and time invariant observed factors explain *none* of the persistence in weight status for primary school-aged children. Moreover, the estimates of γ above 0.9 indicate a substantial degree of persistence. Thus, while persistence from one period to the next is extreme, this persistence is not attributable to or explained by characteristics typically observed by policymakers or health practitioners. Moving to the specification in (3), which replaces the time invariant observed

factors with child-level fixed effects and thereby controls for all time invariant attributes of the child, the estimate of γ falls to 0.775, a decline of roughly 17% from 0.93. This implies that time invariant, *unobserved* factors explain about 17% of the observed persistence in weight z -scores. Examples of such factors include genetic endowments, prior health shocks determined *in utero* or during infancy, time invariant environmental factors such as the presence of grocery stores or outdoor amenities, etc.

When we divide the sample into different sub-groups, we find that the results are predominantly unchanged in the specifications omitting the fixed effects. The only minor difference we see is a slightly higher level of persistence for males relative to females (approximately 0.95 to 0.91, statistically significant at the $p < 0.01$ confidence level). However, once we include child-level fixed effects, the results vary in several cases. For whites, we find that time invariant, unobserved factors explain roughly 26% of overall persistence; only about 4% for non-whites. For males, the fixed effects explain over 70% of overall persistence as the estimate of γ falls to 0.276 (standard error is 0.056). For females, the point estimate for γ increases well above unity and is relatively imprecise. When splitting the sample by mother’s education, we find that time invariant, unobserved factors explain only 5% of total persistence for children with a college educated mother, but roughly 20% for those without a mother without a four-year college degree. Similarly, we find that the fixed effects explain about 4% of total persistence for urban residents, but roughly 23% for non-urban residents. Finally, we obtain little difference across groups when dividing the sample by SES status.

To interpret these findings, it is important to remember that the decline in γ when conditioning on the fixed effects represents the amount of persistence due to time invariant unobserved risk factors. Consequently, we find that overall persistence is fairly extreme as a one standard deviation increase in weight is associated with roughly a 0.9 standard deviation increase in the subsequent period. However, time-varying and time invariant observed attributes explain none of this persistence. Moreover, time invariant unobserved factors also explain very little of the persistence (typically less than one-third). Thus, much of the persistence in child weight is attributable to state dependence, which implies that early interventions that are successful in reducing child weight will have long-run effects. Unfortunately, since our covariates explain little of the variation in weight, identifying such early interventions may be difficult.¹²

Table 2 displays the simulation results for the $\Pr(y_{iT} \geq 85^{\text{th}} \text{ percentile} \mid y_{i0} \geq 85^{\text{th}} \text{ percentile})$. For the full sample, the benchmark probability is 0.84. In other words, in our sample, 84% of children above the 85th percentile in the initial period remain above the 85th percentile in the terminal period. This is consistent with a high degree of persistence in weight. To explore the sources of this persistence, we turn to the simulations.

Panel I contains the simulated probabilities when time-varying unobservables are ignored (i.e., $\varepsilon_{it} = 0$ for all i, t) and time invariant heterogeneity is altered first by removing it entirely (by setting α at the sample mean of $\hat{\alpha}$) and then by retaining the heterogeneity in α , but breaking its correlation with x and y_0 by giving each child a random draw from the empirical distribution of $\hat{\alpha}$. In the first case, the conditional probability of staying above the 85th

¹²The full set of results are available upon request. While some estimated coefficients are statistically significant at conventional levels, the magnitudes are quite small; even the long-run effects of permanent changes in the covariates, given by $\beta/(1 - \gamma)$, are quite small. That said, while our covariate set does include a wide array of the usual family background variables, we do not have information on many recent interventions designed to combat obesity, such as education efforts, healthy food programs, and efforts to promote physical activity. We also do not have data on parents’ height or weight. We return to the issue of parental anthropometric status later.

percentile falls to about 0.753; it falls to roughly 0.576 in the second case. The fact that the conditional staying probability drops noticeably from the benchmark in the second case, but only marginally in the first case, indicates that it is not the *variation* in α across children that determines persistence, but rather the *correlation* between α and the time-varying covariates that explain a little over 30% of total persistence (i.e., $1 - (0.576/0.84)$). Moreover, since the prior results in Table 1 indicate that the time-varying, observed covariates, x , have little explanatory power, this suggests it is really the correlation between α and the initial condition, y_0 , that explains nearly one-third of the total persistence. In other words, children with high initial conditions – measured by weight z -scores upon kindergarten entry – also have high values of α , and this combination is responsible for one-third of the conditional staying probability over the span of kindergarten through eighth grade.

Panels II and III in Table 2 assess whether the importance of α is driven by time invariant observed factors, w , or unobserved factors, η . The first simulation in Panel II sets η equal to zero and leaves w at its actual value. The result is very similar to the first case in Panel I, when α is set equal to its sample mean. In this case, the conditional staying probability is 0.727, implying that it is the setting of η to its sample mean that is driving the first result in Panel I. When instead children are given a random draw for w from its empirical distribution, the probability changes only modestly to 0.703. Again, this is consistent with the results in Table 1 where we found little explanatory power for the time invariant, observed covariates. In Panel III, however, when children retain their own observed factors, x and w , but receive a random draw for η from its empirical distribution, the conditional staying probability falls to 0.593. As such, it is the correlation between time invariant, *unobserved* factors and the initial condition, y_0 , that is responsible for roughly one-third of the conditional staying probability. In other words, children with high initial conditions also have high values of η , and this combination is responsible for one-third of the persistence in weight from kindergarten through eighth grade.

Lastly, Panels IV, V, and VI report the simulated probabilities obtained when children retain their α , but receive draws of either time-varying, observed, x , or unobserved, ε , attributes or both from their respective empirical distributions. The results indicate no impact from altering either, again consistent with the the prior results in Table 1. In sum, the simulations for the full sample indicate that about one-third of the conditional staying probability for weight is due to *persistent, unobserved* risk factors such as genetic endowments, early life health shocks, time invariant environmental factors, etc. The remainder is due to state dependence. The fact that two-thirds of persistence is due to state dependence is encouraging in that early interventions, to the extent that they are successful in reducing weight prior to kindergarten, can have long-run effects on weight during middle school.

The remainder of Table 2 reports the simulated probabilities for the different sub-groups. In addition to the simulations just discussed for the full sample, additional simulations are conducted. Specifically, when drawing from the empirical distributions, we draw not only from the full sample, but also from within one’s own group and outside one’s own group. This enables us to see the effects of differences in the distributions of the various components of the model across groups.

In the interest of brevity, we highlight a few salient findings. First, the benchmark probabilities differ little by gender or urban status. However, non-white children, children with a mother without a four-year college degree, or children residing in a low SES household have a higher benchmark conditional staying probability (race: 0.861

versus 0.823; education: 0.870 versus 0.748; SES: 0.880 versus 0.820). Second, as in the full sample, altering values for the time-varying, observed and/or unobserved factors, x and ε , has little impact on persistence in weight for all demographic groups.

Third, altering values for α , or its components, matters across all demographic groups, but in different ways. For non-whites, children with a mother without a four-year college degree, and children residing in a low SES household, replacing α with the (full) sample mean has little effect on the conditional staying probability. This suggests that these groups have such poor initial conditions, y_0 , that even replacing α with the sample mean is not sufficient to move children in these groups who are initially above the 85th percentile below the 85th percentile in the terminal period. Instead, only when α is replaced by a random draw, particularly a random draw from outside one’s own group, does the conditional staying probability drop to 0.50-0.60. Fourth, while the distributions of α do not differ by much across the different groups, the distributions of the observed component, w , does. In particular, females, children in urban residences, children with a four-year college educated mother, and children in high SES families possess time invariant, observed factors associated with persistence. However, the fact that the overall distribution of α differs little across groups indicates that much of the variation in α is due to the unobserved component, η , which differs little across groups. Thus, in the end, the amount of persistence due to the fixed effects as opposed to state dependence is roughly constant across the groups.

Before turning to the next table, note that the preceding simulation results (and more) can be gleaned from plots of the various distributions provided in Figures B1 and B2 in Appendix B. Figure B1 plots the overall distributions of weight z -scores by demographic group in column one, the distributions of $x\hat{\beta}$ in column two, the distributions of $\hat{\alpha}$ in column 3, and the distributions of $\hat{\varepsilon}$ in column four. When viewing the figures, it is important to pay particular attention to not only *differences* in the distributions across groups, but also the *scale* of the horizontal axis. For example, in the top row, while the distribution of time-varying, observed attributes, $x\hat{\beta}$, is quite different across racial groups, the distributions are concentrated over a range of -0.1 to 0.1; the overall distributions of weight z -scores range from about -2 to 2. Thus, while time-varying, observed covariates differ across racial groups, they explain little of the overall variation in weight. In contrast, the distributions of time-varying, unobserved attributes, $\hat{\varepsilon}$, exhibit meaningful variation overall, but the distributions are virtually identical across demographic groups.

Figure B2 reproduces the distributions of $\hat{\alpha}$ in column 1 along with its decomposition into observed attributes, $w\hat{\delta}$, in column 2 and unobserved attributes, $\hat{\eta}$, in column 3. As in Figure B1, we see that while the distributions of observed characteristics differ, for example, along racial lines in the top row, the scaling is such that the distributions of $\hat{\alpha}$ are determined predominantly by the distributions of $\hat{\eta}$ which differ little between whites and non-whites. In sum, then, the figures indicate little variation across demographic groups in terms of the overall distribution of weight z -scores or the components most responsible for variation in weight.

Table 3 displays the analogous results for the $\Pr(y_{iT} \geq 95^{\text{th}} \text{ percentile} \mid y_{i0} \geq 95^{\text{th}} \text{ percentile})$. Compared to the results in Table 2, three primary differences emerge. First, the benchmark probability is lower in the full sample and for each demographic group (e.g., 0.762 for the full sample). Thus, there is less persistence in the extreme upper tail of the weight distribution. Moreover, the difference in the benchmark probability across each demographic group is now economically meaningful (race: 0.732 versus 0.795 favoring whites; gender: 0.710 versus 0.807 favoring

females; urban status: 0.749 versus 0.791 favoring non-urban; education: 0.646 versus 0.790 favoring four-year college educated; SES: 0.740 versus 0.799 favoring high SES). Second, the vast majority of the persistence is due to time invariant heterogeneity, α ; even more so than in Table 2. State dependence, as well as time-varying factors, x and ε , do not play much of a role in explaining persistence in the extreme upper tail. For example, replacing α with the sample mean for all children reduces the conditional staying probability in the full sample to less than 15% and less than 20% within each demographic group. Even replacing α with a random draw from its empirical distribution cuts the conditional staying probability by nearly one-half in all cases. Third, unlike in Table 2, we find that setting η to zero in Panel II results in lower conditional staying probabilities than in Panel III when η is replaced by random draws from different empirical distributions. This indicates that giving children initially above the 95th percentile an average draw from the distribution of η (i.e., setting η to zero) is sufficient to bump most of these children below the 95th percentile by the terminal period, whereas this is not sufficient when using the 85th percentile as the threshold.

Finally, Table 4 presents the results for the $\Pr(y_{iT} \geq 85^{\text{th}} \text{ percentile} \mid y_{i0} \leq 50^{\text{th}} \text{ percentile})$. This case illuminates factors associated with relatively extreme weight gain during early childhood (i.e., sizeable upward mobility as opposed to persistence). In terms of the benchmark case, the probability of moving from below the median at kindergarten entry to above the 85th percentile by the end of eighth grade is roughly 12% in the full sample. While this probability does not differ much across the demographic groups, small differences arise favoring non-whites, urban residents, and children with a mother with a four-year college degree and those residing in high SES households (race: 0.113 versus 0.121; gender: 0.104 versus 0.132; urban: 0.104 versus 0.124; education: 0.079 versus 0.131; SES: 0.106 versus 0.145).

Turning to the simulations, we obtain a few noteworthy findings. First, time-varying factors, x and ε , continue to not play any meaningful role. Second, replacing α with the sample mean reduces the probability of crossing the 85th percentile conditional on starting below the median to zero in all cases. Replacing α with a random draw from different empirical distributions roughly doubles the probability of crossing the 85th percentile relative to the benchmark in all cases. Together, these results imply that children initially below the median tend to have favorable values of α . Specifically, α is not randomly distributed in the population, but rather has a positive (partial) correlation with the initial condition, y_0 . Only the few children with extremely unfavorable draws of α , despite being below the median in the initial period, experience extreme upward mobility. Moreover, if α were randomly assigned, the probability of moving from below the median to above the 85th percentile would roughly double. This is a testament to the importance of time invariant factors, α , in determining weight status.

Third, the effect of randomly assigning α is due to randomly assigning time invariant, unobserved factors, η . Randomly assigning the time invariant, observed factors, w , has little impact on the probability of extreme upward mobility. Moreover, removing time invariant, unobserved factors by setting η to zero reduces the probability of extreme upward mobility to nearly zero in all cases. The implication is that children below the median tend to have favorable draws of α , which really means favorable draws of time invariant, unobserved factors, η .

4.2 Height

Next we turn to the analysis of height. While height *per se* is not a policy concern in the U.S., it is interesting to compare the dynamics of height with those of weight. In addition, it is useful to examine the individual components

of BMI prior to assessing BMI z -scores in the next section.

Table 5 displays the results for height z -scores. For the full sample, the estimates of γ across the first two specifications are very similar to those using weight z -scores; namely, 0.937 and 0.936 (standard errors are 0.004 and 0.004, respectively). However, the estimate of γ falls to 0.603 (standard error is 0.048) in the fixed effect specification (compared to 0.775 in Table 1). As in Table 1, the estimate of γ is statistically significant at the $p < 0.01$ confidence level, all three specifications are strongly identified, the estimate of γ barely changes when we include time-varying and time invariant observed attributes, and the estimates of γ above 0.9 in the first two specifications indicate a substantial degree of persistence. Thus, as in Table 1, while anthropometric measures are quite persistent from one period to the next, this is not attributable to or explained by observed characteristics.

In contrast to weight z -scores, the child-level fixed effects explain about 36% of the overall persistence in child height (versus only 17% for weight z -scores). This is perhaps not surprising as unobserved biological factors – most noticeably, parental height – are not included in our set of observed covariates. The fact that time invariant, unobserved attributes account for a greater share of the persistence in height implies that state dependence, and thus the long-run impact of successful, early interventions – that do not alter relevant, time invariant, unobserved attributes – is diminished. For example, a *one-time* intervention that reduces a child’s weight by one standard deviation prior to kindergarten entry, *ceteris paribus*, is expected to reduce the child’s weight by over one-third of a standard deviation in spring eighth grade. Thus, one-third of the effects of the early intervention persist through eighth grade. An intervention that raises a child’s height by one standard deviation prior to kindergarten entry, *ceteris paribus*, is expected to increase the child’s height only by slightly over 0.10 standard deviations in spring eighth grade. As such, only about one-tenth of the effects of the early intervention persist through eighth grade; the remainder of the intervention dies out.

When we divide the sample into different sub-groups, we find that the results are predominantly unchanged in the specifications omitting the fixed effects. The only minor difference we see is a slightly higher level of persistence for males relative to females and non-urban residents relative to urban residents (approximately 0.95 to 0.92 and statistically significant at the $p < 0.01$ confidence level in each case). However, as with weight z -scores, once we include child-level fixed effects, the results vary in several cases. When we split the sample by race, we find that time invariant, unobserved factors explain roughly 41% of overall persistence for whites versus about 28% for non-whites. For males, the fixed effects explain over 50% of overall persistence as the estimate of γ falls to 0.460 (standard error is 0.055). For females, the point estimate falls to 0.739 (standard error is 0.079); thus, accounting for only about 20% of overall persistence. When we divide the sample by mother’s education, we find that time invariant, unobserved factors also explain over 50% of total persistence for children with a college educated mother; roughly 30% for those with a mother without a four-year college degree. Similarly, we find that the fixed effects explain about 40% of total persistence for children in high SES households, but roughly 25% for children in low SES households. Finally, we obtain little difference across groups when dividing the sample by urban status.

Tables 6-8 present the analogous set of simulation results for height z -scores; Figures B3 and B4 display the plots. We discuss the results briefly. In terms of the benchmark probabilities, a few differences emerge relative to the previous results for weight. First, the benchmark probabilities are lower for height than the corresponding

probabilities for weight in all cases across Tables 6-7. For example, $\Pr(y_{iT} \geq 85^{\text{th}} \text{ percentile} \mid y_{i0} \geq 85^{\text{th}} \text{ percentile})$ and $\Pr(y_{iT} \geq 95^{\text{th}} \text{ percentile} \mid y_{i0} \geq 95^{\text{th}} \text{ percentile})$ are 0.606 and 0.467, respectively, in the full sample for height; 0.840 and 0.762, respectively, for weight. Thus, persistence in the upper half of the distribution is lower, albeit still high, for height. Second, while there may exist more mobility in terms of height, extreme upward mobility for height is less common than for weight. In the full sample, $\Pr(y_{iT} \geq 85^{\text{th}} \text{ percentile} \mid y_{i0} \leq 50^{\text{th}} \text{ percentile})$ is 0.030 for height and 0.118 for weight.

Turning to the simulations, a few patterns emerge. First, while the time-varying factors, x and ε , have a bit more impact on height than weight, their combined effect is still modest. In Tables 6-8, replacing x and/or ε with different values increases the conditional staying probabilities in all cases for the full sample. This indicates that, on average, children initially above the median tend to have less favorable (in terms of raising height) time-varying attributes, partially offsetting the child’s height in the initial period.

Second, as with weight, most of persistence in height is attributable to time invariant factors captured by α . However, the patterns are different. In Tables 6 and 7, we find that replacing α with the sample mean drops the conditional staying probabilities above the 85th and 95th probabilities to zero for the full sample and all demographic groups. Further analysis reveals that this stems from the unobserved component captured by η ; varying the time invariant, observed component, w , has little effect. This implies that children in the upper tail of the height distribution upon entry to kindergarten possess time invariant, unobserved attributes that tend to keep them in the upper tail. Replacing these attributes with the sample mean, or a random draw, essentially guarantees these children will fall out of the upper tail by the end of eighth grade. Replacing the unobserved component of the fixed effects, η , with a random draw similarly reduces the conditional staying probabilities, but not as much; the probabilities fall to around 0.25 and 0.10 in Tables 6 and 7, respectively. This is perhaps not surprising as genetics and early biological factors presumably play a large role in determining child height.

Third, Table 8 suggests that extreme upward mobility in height is rare since children initially below the median have unfavorable draws of time invariant, unobserved heterogeneity, η . Replacing η with its sample average would eliminate extreme upward mobility entirely as the few cases of observed extreme upward mobility is due to a handful of children having very favorable values of η despite being below the median upon entry to kindergarten. On the other hand, replacing η with a random draw would increase extreme upward mobility by four- to five-fold.

Finally, Figures B3 and B4 provide a graphical representation of these findings. The implications are very similar to those discussed above with respect to the figures for weight. The only subtle difference, consistent with the simulation results, is that the distribution of the fixed effects, $\hat{\alpha}$, explain a bit more of the overall variation in height (with this variation reflecting the unobserved component, $\hat{\eta}$).

4.3 BMI

Next we turn to the analysis of BMI. Table 9 presents the regression results. For the full sample, the estimates of γ across the first two specifications are very similar to those in Tables 1 and 5; namely, 0.912 and 0.911 (standard errors are 0.004 and 0.005, respectively). However, the estimate of γ now falls to 0.217 (standard error is 0.015) in the fixed effect specification (compared to 0.775 and 0.603 in Tables 1 and 5, respectively). As in Tables 1 and

5, the estimate of γ is statistically significant at the $p < 0.01$ confidence level, all three specifications are strongly identified, the estimate of γ barely changes when we include time-varying and time invariant observed attributes, and the estimates of γ above 0.9 in the first two specifications indicate a substantial degree of persistence. Thus, as with weight and height z -scores, while persistence from one period to the next in BMI z -scores is high, it is not attributable to or explained by observed characteristics.

While the first two specifications differ little across Tables 1, 5, and 9, the results from the fixed effect specification do. As noted above, time invariant, unobserved factors account for roughly 17% of the total persistence in weight z -scores and 36% for height z -scores. For BMI, the fixed effects now account for nearly 80% of total persistence. The economically and statistically meaningful drop in the estimate of γ implies a substantially smaller role for state dependence in the persistence of child BMI. Consequently, the long-run impact of early interventions – that do not alter relevant, time invariant, unobserved attributes – on BMI is quite small. For example, a *one-time* intervention that reduces a child’s BMI prior to kindergarten entry by one standard deviation, *ceteris paribus*, is expected to have essentially no impact on BMI in spring eighth grade. A *permanent* intervention that reduces a child’s BMI by 0.10 standard deviations *every period*, will only result in a long-run decrease in the child’s BMI of roughly 0.13 standard deviations. This has profound implications for the types of policies one should pursue if the objective is to reverse the obesity epidemic.

When we divide the sample into different sub-groups, we find that the results are qualitatively similar across all demographic groups for each of the three specifications, in contrast to the prior results for weight and height. In terms of the first two specifications, there are essentially no differences across the various groups. For the fixed effect specification, the only minor difference of note is for gender. In this case, the fixed effects account for approximately 80% of total persistence for males and roughly 70% for females. For all the remaining divisions of the sample, time invariant factors account for roughly 73 - 78% of total persistence. Again, this is a striking finding as it indicates that while there may be *level* differences in BMI across demographic groups, the extent and origins of *persistence* are not fundamentally different across groups.

Tables 10-12 display the simulation results for BMI z -scores; Figures B5 and B6 contain the plots. In Tables 10 and 11, the benchmark probabilities lie in between the conditional staying probabilities for weight and height reported in the corresponding Tables 2-3 and 6-7. This is also true for most of the demographic sub-groups. Furthermore, the benchmark probabilities are consistent with the high degree of persistence in BMI documented earlier. For example, the conditional probability of staying above the 85th percentile is 0.746 in the full sample (see Table 10); 0.715 for staying above the 95th percentile (see Table 11). Lastly, the benchmark probabilities are notable in that the gaps between racial, education, and SES groups in Tables 10 and 11 are larger than the corresponding gaps for either weight or height separately. For instance, the conditional probability of staying above the 95th percentile for BMI is 0.664 for whites and 0.769 for non-whites. The corresponding gap for weight (height) is 0.732 versus 0.795 (0.481 versus 0.446). Thus, demographic differences in persistence of remaining in the upper tail of the BMI distribution are sizeable.

When we turn to the simulated probabilities, a few findings stand out. First, altering the values of the time invariant components in Panels I, II, and III of Tables 10-11 yields results that are qualitatively similar to those

reported in Tables 6-7 for height. In particular, in Panel I we find that replacing α with the sample mean reduces the conditional probability of staying above the 85th and 95th percentiles to zero in nearly every case. Moreover, this is predominantly due to the salient role of time invariant, unobserved factors, η . Variation in time invariant, observed factors, w , explain a modest amount of variation in the conditional probability of staying above the 85th percentile (see Table 10), but not when using the 95th percentile as the threshold (see Table 11). Thus, the results are consistent with children in the upper part of the BMI distribution possessing less favorable time invariant factors, particularly those unobserved. The results are also consistent with Figures B5 and B6 which indicate that the majority of the variation in BMI is due to the fixed effects, $\hat{\alpha}$, and the unobserved component, $\hat{\eta}$, in particular.

Second, in Panel II of Table 10, where variation in time invariant, observed factors, w , plays a modest role, we find that whites, females, non-urban residents, children with a mother with a four-year college degree, and children in high SES households continue to possess more favorable attributes. The largest discrepancy occurs along racial lines. If we set η to zero and give white children a random draw of w from the empirical distribution for whites (non-whites), we obtain a conditional staying probability of 0.015 (0.121). Setting η to zero and giving non-white children a random draw of w from the empirical distribution for whites (non-whites), we obtain a conditional staying probability of 0.015 (0.118). Thus, the variation in the distribution of time invariant, observed factors is responsible for roughly a ten percentage point difference along racial lines in the conditional probability of remaining above the 85th percentile, *ceteris paribus*. Finally, as in all the analysis of weight and height, we find very little role for variation in time-varying factors, either observed or unobserved.

Table 12 presents the results for the $\Pr(y_{iT} \geq 85^{\text{th}} \text{ percentile} \mid y_{i0} \leq 50^{\text{th}} \text{ percentile})$. In terms of the benchmark probabilities for extreme upward mobility, we obtain higher probabilities for BMI than either weight or height. For example, the probability of having a BMI above the 85th percentile in the terminal period conditional on entering kindergarten below the median is 0.142 for the full sample (see Table 12). The corresponding figures are 0.118 and 0.003 for weight and height, respectively. The probability of extreme upward mobility for BMI is particularly high for whites, children with non-college educated mothers, and children residing in low SES households (0.167, 0.162, and 0.192, respectively).

Turning to the simulations, we obtain a few findings. First, time-varying factors, x and ε , continue to not play any meaningful role. Second, replacing α with the sample mean reduces the probability of crossing the 85th percentile conditional on starting below the median to zero in all cases, just as in Tables 4 and 8. Replacing α with a random draw from different empirical distributions roughly increases the probability of crossing the 85th percentile by two- to three-fold relative to the benchmark in all cases. Together, these results continue to imply that children initially below the median tend to have favorable values of α . Only a few children with extremely unfavorable draws of α , despite being initially below the median, experience extreme upward mobility. Moreover, if α were randomly assigned, the probability of moving from below the median to above the 85th percentile would increase substantially.

Third, the effect of altering α is due to altering the time invariant, unobserved factors, η . However, as in Table 10, the time invariant, observed factors, w , explain a modest amount of the variation in the probability of extreme upward mobility overall, as well as across racial, education, and SES groups. Specifically, whereas removing time invariant, unobserved factors by setting η to zero reduces the probability of extreme upward mobility to nearly zero

for weight and height, this is not the case for BMI as the probability varies from roughly 1 - 8%.

4.4 Discussion

While there are many subtle results emerging from the analysis, perhaps the most important is that persistence in weight, height, and BMI is quite high over the period spanning kindergarten through eighth grade and that this persistence is predominantly driven by *persistent, unobserved* heterogeneity. Time-varying observed and unobserved factors play little role. Time invariant, *observed* heterogeneity plays a modest role in some instances. In particular, children who are male or black, rural or northeast residents, non-native English speakers, had a high birthweight, and have a mother with low education, a low age at first birth, or who participated in the labor force during the child’s infancy tend to have higher BMI (as evidenced by inspection of the estimation results of (6)). State dependence plays a prominent role for weight only. That said, it is worth re-iterating that the majority of persistence in weight, height, and BMI is due to time invariant, unobserved factors.

This finding implies that, while earlier intervention is preferred to later interventions, *only* interventions that alter the crucial, time invariant, unobserved risk factors captured by η are likely to be effective in the long-run. Interventions that leave the attributes captured by η unaltered are likely to have, at best, minimal short-run effects and little to no long-run effects. This is entirely consistent with the findings reported in Davis and Gebremariam (2010). There, the authors document that community-based interventions designed to combat childhood obesity that were deemed as successful according to the analysis of data collected via randomized control trials did not produce lasting effects. Eventually, children returned to their “natural state” (p. 22).

This naturally begs the question concerning the attributes reflected by η . From the analysis presented here, all we can conclude is that they are not contained in our set of covariates taken from the ECLS-K and they do not vary during the primary school years. The prior literature, discussed earlier, posits some possibilities: prenatal attributes such as maternal BMI, maternal weight gain, maternal smoking, and gestational diabetes requiring insulin and post-natal attributes such as breastfeeding, transitions to solid foods, and age at adiposity rebound. While we do control for birthweight, birthweight alone is not a sufficient proxy for these early influences on fetal development as noted earlier. Finally, while time invariant, environmental factors, such as neighborhood characteristics, are also captured by η , prior evidence suggests that these are not likely to play a significant role. For example, prior studies using twins that are reared apart conclude that familial environment does not play a salient role (Eriksson et al. 2001). In an attempt to delve further into this issue, we undertake one final analysis using the ECLS-B. We turn to it now.

5 ECLS-B

5.1 Data

To explore the early life origins of anthropometric persistence, we utilize data from the restricted version of the ECLS-B. Collected by the US Department of Education, the ECLS-B collects information on a nationally representative

cohort of children born in 2001 at 9 months of age, two years, four years, and five years. As with the ECLS-K, our final sample consists of a balanced sample of children for whom we have valid measures of age, gender, height, and weight.¹³ Given the age of the sample, we convert weight into z -scores; height is measured in centimeters.

The following time invariant covariates are included: gender, race/ethnicity (white, black, Hispanic, Asian, and other), mother’s age at first birth, birthweight indicators (normal or low), indicator for intrauterine growth retardation (less than 10%, 10-24%, 25-49%, 50-75%, 76-89%, and 90% and above)¹⁴, indicator for premature birth, indicator for birth status (singleton, twin, or higher order birth), mother’s height, mother’s weight prior to pregnancy, mother’s weight gain during pregnancy, indicator for prenatal care (inadequate, intermediate, adequate, or adequate plus), indicator for maternal prenatal vitamin consumption within the three months preceding conception, indicator for maternal prenatal vitamin consumption during the first trimester, indicator for maternal smoking within the three months preceding conception, indicator for maternal smoking within the third trimester, indicator if mother has smoked more than 100 cigarettes in her lifetime, indicator for maternal alcohol consumption within the three months preceding conception, number of current smokers in the household, region (northeast, midwest, south, and west), city type (urban, suburban, or rural), indicator for mother’s participation in WIC during pregnancy, indicator for mother’s participation in WIC during infancy, and scores on infant mental and motor assessments administered at 9 months.

The following time-varying covariates are included: age, mother’s education (less than high school, high school/GED, some college, four-year college degree, and more than four years of college), an index of SES status, indicator for the household being in poverty, number of children’s books in the household, household size, family type (two parents plus siblings, two parents and no siblings, one parent and siblings, one parent and no siblings, and other), indicator for biological mother present, indicator for biological father present, indicator for no father present, indicator for no mother present, indicator for parental respondent’s marital status, indicator of current TANF participation, indicator of current FSP participation, indicator for health insurance, indicator for current medicaid participation, indicator for current WIC participation, indicator for household food security (household never worried about running out of food), hours per day spent watching television during the school week, indicator for household rules regarding television watching, neighborhood safety (very safe, somewhat safe, and not safe), mother’s labor force status (full-time, part-time, and not working), indicators for primary child care arrangement (parents, other relatives, non-relatives, center-based care, or Head Start), indicator for school enrollment, indicator if English is the primary home language, and mother’s weight. For all covariates (except gender, age, height, and weight), we include dummy variables for missing observations.

5.2 Results

The results are presented in Tables 13-20. Tables 13 and 17 display the regression estimates; the remaining tables present the simulation results. Figures B7-B10 in Appendix B contain the corresponding plots.

¹³The possible sample size is roughly 6,950; the initial sample size in the first wave is about 10,700. Restricting the sample to those with valid data on age, gender, height, and weight reduces the sample size to approximately 5,450. This is the final sample size per wave in the regression analysis. Note, all sample sizes are rounded to the nearest 50 per NCES restricted data regulations for the ECLS-B.

¹⁴Intrauterine growth retardation measures the ratio of birthweight to predicted weight based on gestation age.

In terms of the coefficient estimates, the results in Table 13 using weight z -scores are fairly similar to those obtained using the ECLS-K when omitting child-specific fixed effects. Specifically, the estimates of γ in the full sample and each of the demographic sub-groups is statistically significant and ranges from 0.84 to 0.89. As with the ECLS-K, the fact that the estimate of γ does not change moving from (1) to (2) implies that our lengthy vector of time-varying and time invariant observed factors explain none of the persistence in weight status for infants and young children. Given the additional time invariant controls available in the ECLS-B, this is striking. Moreover, the estimates of γ near 0.9 indicate a substantial degree of persistence even prior to kindergarten. However, unlike in the ECLS-K, inclusion of child-level fixed effects explains the majority of this persistence. Here, the estimate of γ falls to 0.124 (standard error of 0.013) in the full sample; the estimates vary from 0.105 to 0.144 across the various sub-groups. This implies that *time invariant, unobserved* factors explain about 85% of the observed persistence in weight z -scores during early childhood. In contrast, only 17% of observed persistence in weight z -scores during primary school is due to time invariant, unobserved heterogeneity. Again, given that we observe many more time invariant attributes of children in the ECLS-B, this is a startling result.

Table 17 displays the corresponding regression results for height. Four interesting patterns emerge. First, persistence in height in the models not controlling for any other covariates – based on the specification in (1) – is of a much smaller magnitude than found in the ECLS-K when assessing height for older children or in the ECLS-B when assessing weight. Second, when controlling for observed heterogeneity – based on the specification in (2) – persistence actually increases by about 15%. This is consistent with a negative correlation between the initial condition for height, y_0 , which is really ‘length’ at nine months of age, and observed heterogeneity associated with greater height. Third, as with weight in the ECLS-B, there is very little difference across the demographic sub-groups. Finally, when child-level fixed effects are included, the estimates of γ become *negative* and statistically significant (although always below 0.06 in absolute value). Thus, *all* of the persistence in child height up to age five is attributable to *time invariant, unobserved* heterogeneity.

Tables 14-16 and 18-20 report the results of the same simulations performed as when using the ECLS-K. In the interest of brevity, we only briefly summarize the results. First, time-varying attributes, both observed and unobserved have no effect on persistence. Given the lengthy vector of attributes, as well as the plethora of time-varying, unobserved attributes captured by ε , this continues to be a noteworthy finding. Second, the benchmark probabilities for both weight and height, along with the distributions of observed and unobserved, time-varying attributes, do not qualitatively differ across the demographic sub-groups (see also Figures B7 and B9). As such, not only do the benchmark probabilities not vary across groups, but replacing a child’s own x and/or ε with draws from the opposite group has no discernible effect on persistence.

Third, as in the ECLS-K, time invariant heterogeneity continues to play a prominent role in understanding persistence in child weight and height. For weight, replacing α with its sample mean explains virtually all persistence through age five. Moreover, replacing the fixed effect of a child initially below the median with the sample mean roughly doubles the probability that the child’s weight will exceed the 85th percentile at age five. For height, replacing α with its sample mean explains most, but not all, persistence. However, replacing the fixed effect of a child initially below the median with the sample mean does not alter the probability that the child’s height will exceed the 85th

percentile at age five.

Fourth, time invariant, *observed* attributes play a more prominent role, particularly for height, in explaining persistence up to age five than in the ECLS-K analysis of primary school children. This could be attributable to two sources. On the one hand, the time invariant, observed attributes may play a more important role in the determination of child weight and height prior to age five. On the other hand, the vector of controls is not identical across the two data sources. Examining the results of (6), the most important covariates relate to birthweight, birth status (i.e., singleton, twin, or higher order birth), intrauterine growth retardation, breastfeeding duration, mother’s height, and mother’s weight gain during pregnancy. That said, as measured by the R^2 , only 19% (22%) of the variation in $\hat{\alpha}$ is explained by the covariates included in (6) when examining weight (height).

Finally, time invariant attributes, both observed and unobserved, differ across the various demographic subgroups, particularly along racial lines. For example, in Table 14, the probability of a white (non-white) child’s weight persisting above the 85th percentile when the child’s own fixed effect, α , is replaced by a random draw from the sample distribution is 0.284 (0.242). Replacing the child’s own fixed effect, α , with a random draw from the sample distribution for the child’s own racial group, the probability of persisting above the 85th percentile is 0.216 (0.284). Replacing the child’s own fixed effect, α , with a random draw from the sample distribution from the opposite racial group, the probability of persisting above the 85th percentile is 0.333 (0.179). Similar patterns hold in the other panels for weight and height.

In sum, the results from the sample of children aged five and younger in the ECLS-B are consistent with the sample of primary school children in the ECLS-K. Namely, persistence in weight and height is quite high, and this persistence is mainly driven by *time invariant* heterogeneity. However, in contrast to the primary school sample, time invariant, *observed* attributes play a bit more of an important role. In particular, while the associations between birthweight, gestation age, maternal height and weight, and single versus multiple birth and fetal development are not strong, perhaps due to the complexities involved these relationships that are only currently beginning to be understood in the medical literature, these controls do play a small role in explaining persistence. Nonetheless, the primary determinants of fetal and infant development that may be critical in placing children on an “obesity trajectory” remain unobserved, even in the ECLS-B. Such unobserved attributes are likely to include gestational diabetes treated with insulin or periods of undernutrition during pregnancy. Similarly, recent research has uncovered genetic abnormalities associated with obesity (e.g., Asai et al. 2013; Ramachandrappa et al. 2013).

6 Conclusion

Concern over childhood and adult obesity has risen dramatically over the past decade. As this concern has risen, our understanding that interventions earlier in life are likely to have greater impact has risen as well. This understanding follows from well chronicled evidence that obesity is highly persistent; as such, adults may become trapped on an “obesity trajectory” early in life. However, little is known about the origins of these correlations in anthropometric measures over the life cycle. Specifically, whether this correlation reflects state dependence, observed heterogeneity, or unobserved heterogeneity is unknown. Moreover, when this persistence in weight status begins – adolescence,

early childhood, postnatally, or prenatally – is also unknown. Prior work has identified three ‘critical’ development periods as it relates to obesity: *in utero*, adiposity rebound (around age four to six), and adolescence. However, as Dietz (1997, p. 1886S) notes, “The relative contribution of each of these critical periods to the prevalence, morbidity and mortality of adult obesity remains uncertain.”

Better understanding of the dynamics of weight status is crucial for sound policymaking. If weight is highly persistent and the source of this persistence is state dependence, then small (permanent) changes will have large, long-run effects even if the contemporaneous effects are small. However, if persistence is due to biological or environmental factors that are time invariant, then the *only* changes that will have long-run effects are those that alter these underlying factors. Absent such effects, interventions will not alter the long-run anthropometric status of individuals even if they have contemporaneous effects.

The evidence presented here indicates, first, that there is significant persistence in weight and height starting during infancy and, second, that this persistence is predominantly due to time invariant heterogeneity across individuals determined at birth or shortly thereafter, not state dependence. Moreover, little variation in this time invariant heterogeneity is explained by attributes observed in the data analyzed here. The few time invariant, observed attributes that do seem to play a role in the persistence of weight status over the early part of the life cycle relate to fetal and infant nutrition. This suggests that of the three ‘critical’ periods noted in Dietz (1997), *in utero* (and post-natal) plays the largest role. It also suggests that strategies to reverse the current childhood obesity epidemic may need to start even earlier than previously thought, namely *in utero*. This confirms recent policy prescriptions advocated elsewhere. For example, Brisbois et al. (2012, p. 347) concludes: “Given that obesity may be programmed *in utero* and during early infancy, preventive measures should be initiated preconception, during pregnancy and continue throughout early childhood.” Examples of such measures may include altering institutional rules concerning federal nutrition programs, such as SNAP or WIC, or education provided under these programs, as they relate to pregnant women (e.g., Baum 2012).

References

- [1] Almond, D. and J. Currie (2011), “Killing Me Softly: The Fetal Origins Hypothesis,” *Journal of Economic Perspectives*, 25,153-172.
- [2] Asai, M., S. Ramachandrappa, M. Joachim, Y. Shen, R. Zhang, N. Nuthalapati, V. Ramanathan, D. E. Strohlic, P. Ferket, K. Linhart, C. Ho, T. V. Novoselova, S. Garg, M. Ridderstrale, C. Marcus, J. N. Hirschhorn, J. M. Keogh, S. O’Rahilly, L. F. Chan, A. J. Clark, I. S. Farooqi, J. A. Majzoub (2013), “Loss of Function of the Melanocortin 2 Receptor Accessory Protein 2 Is Associated with Mammalian Obesity,” *Science*, 341, 275-278.
- [3] Barker, D.J.P. (1997), “Maternal Nutrition, Fetal Nutrition, and Disease in Later Life,” *Nutrition*, 13, 807-813.
- [4] Baum, C.L. (2012), “The Effects of Food Stamps on Weight Gained by Expectant Mothers,” *Journal of Population Economics*, 25, 1307-1340.
- [5] Bhattacharya, D. and B. Mazumder (2011), “A Nonparametric Analysis of Black-White Differences in Intergenerational Mobility in the United States,” *Quantitative Economics*, 2, 335-379.
- [6] Brisbois, T.D., A.P. Farmer. and L.J. McCargar (2012), “Early Markers of Adult Obesity: A Review,” *Obesity Reviews*, 13, 347-367.
- [7] Cawley, J. and C. Meyerhoefer (2012), “The Medical Care Costs of Obesity: An Instrumental Variables Approach,” *Journal of Health Economics*, 31, 219-230.
- [8] Davis, K. and K.K. Christoffel (1994), “Obesity in Preschool and School-Age Children: Treatment Early and Often May Be Best,” *Archives of Pediatric and Adolescent Medicine*, 148, 1257-1261.
- [9] Davis, M.M. and A. Gebremariam (2010), “Economic Long-Term Impacts of Interventions Aimed at Preventing or Reducing Obesity Among Children,” CCR 62, USDA Economic Research Service.
- [10] Dietz, W.H. (1997), “Periods of Risk in Childhood for the Development of Adult Obesity—What Do We Need to Learn?” *Journal of Nutrition*, 127, 1884S-1886S
- [11] Dietz, W.H. and T.N. Robinson (2005), “Overweight Children and Adolescents,” *New England Journal of Medicine*, 352, 2100–2109.
- [12] Dor, A., C. Ferguson, E. Tan, C. Langwith (2010), “A Heavy Burden: The Individual Costs of Being Overweight and Obese in the United States,” Department of Health Policy, George Washington University.
- [13] Eriksson, J., T. Forsén, J. Tuomilehto, C. Osmond, and D. Barker (2001), “Size at Birth, Childhood Growth and Obesity in Adult Life,” *International Journal of Obesity*, 25, 735-740.
- [14] Finkelstein, E. A. , I.C. Fiebelkorn, and G. Wang (2004), “State-Level Estimates of Annual Medical Expenditures Attributable to Obesity,” *Obesity Research*, 12.

- [15] Finkelstein, E.A. and L. Zuckerman (2007), *The Fattening of America: How The Economy Makes Us Fat, If It Matters, and What To Do About It*, John Wylie and Sons, Inc.
- [16] Freedman, D.S., L.K. Khan, W.H. Dietz, S.R. Srinivasan, and G.S. Berenson (2001), "Relationship of Childhood Obesity to Coronary Heart Disease Risk Factors in Adulthood: The Bogalusa Heart Study," *Pediatrics*, 108, 712–718.
- [17] Freedman, D.S., L.K. Khan, M.K. Serdula, W.H. Dietz, S.R. Srinivasan, and G.S. Berenson (2005), "Racial Differences in the Tracking of Childhood BMI to Adulthood," *Obesity Research*, 13, 928–935.
- [18] Frisvold, D.E. and A. Giri (2013), "The Potential of Early Childhood Education as a Successful Obesity Intervention," in V. Brennan, S. Kumanyika, and R. Zambrana (eds.), *Obesity Interventions in Underserved US Populations: Evidence and Directions*, Johns Hopkins University Press.
- [19] Gable, S., J. Britt-Rankin, and J. Krull (2008), "Ecological Predictors and Developmental Outcomes of Persistent Childhood Overweight," Contractor and Cooperator Report No. 42, USDA, ERS.
- [20] Godfrey, K.M. and D.J.P. Barker (2001), "Fetal Programming and Adult Health," *Public Health Nutrition*, 4, 611-624.
- [21] Ham, J.C., D. Iorio, and M. Sovinsky (2013), "Caught in the Bulimic Trap? Persistence and State Dependence of Bulimia Among Young Women," *Journal of Human Resources*, 48, 736-767.
- [22] Halliday, T. (2008), "Heterogeneity, State Dependence and Health," *Econometrics Journal*, 11, 499–516.
- [23] Hsiao, C (1986), *Analysis of Panel Data*, Cambridge University Press.
- [24] Iughetti, L., M. de Simone, A. Verrotti, M.L. Iezzi, B. Predieri, P. Bruzzi, S. Bernasconi, F. Balli, and G. Bedogni (2008), "Thirty-year Persistence of Obesity after Presentation to a Pediatric Obesity Clinic," *Annals of Human Biology*, 35, 439-448
- [25] Kleibergen, F. and R. Paap (2006), "Generalized Reduced Rank Tests Using the Singular Value Decomposition," *Journal of Econometrics*, 133, 97-126.
- [26] Krebs, N.F., and M.S. Jacobson (2003), "Prevention of Pediatric Overweight and Obesity," *Pediatrics*, 112, 424–430.
- [27] Millimet, D.L. and R. Tchernis (2012), "Estimation of Treatment Effects Without an Exclusion Restriction: with an Application to the Analysis of the School Breakfast Program," *Journal of Applied Econometrics*, forthcoming.
- [28] Millimet, D.L. and R. Tchernis (2013), "Anthropometric Mobility During Childhood Obesity," unpublished manuscript, Southern Methodist University.
- [29] Mocan, N. and E. Tekin (2011), "Obesity, Self-Esteem, and Wages," in M. Grossman and N. Mocan (eds.) *Economic Aspects of Obesity*, University of Chicago Press.

- [30] Nader P.R., M. O'Brien, R. Houts, R. Bradley, J. Belsky, R. Crosnoe, S. Friedman, Z. Mei, and E.J. Susman (2007), "Identifying the Risk for Obesity in Early Childhood," *Pediatrics*, 118(3), e594-e601.
- [31] Ogden, C.L., K.M. Flegal, M.D. Carroll, and C.L. Johnson (2002), "Prevalence and Trends in Overweight among US Children and Adolescents, 1999-2000," *Journal of the American Medical Association*, 288(14), 1728-1732.
- [32] Ogden, C.L., M.D. Carroll, and K.M. Flegal, (2008), "High Body Mass Index for Age Among US Children and Adolescents," *Journal of the American Medical Association*, 299, 2401-2405.
- [33] Ramachandrappa, S., A. Raimondo, A.M.G. Cali, J.M. Keogh, E. Henning, S. Saeed, A. Thompson, S. Garg, E.G. Bochukova, S. Brage, V. Trowse, E. Wheeler, A.E. Sullivan, M. Dattani, P.E. Clayton, V.Datta, J.B. Bruning, N.J. Wareham, S. O'Rahilly, D.J. Peet, I. Barroso, M.L. Whitelaw, I.S. Farooqi (2013), "Rare Variants in Single-Minded 1 (SIM1) are Associated with Severe Obesity," *Journal of Clinical Investigation*, 123, 3042-3050.
- [34] Ulrick, S.W. (2008), "Using Semi-parametric Methods in an Analysis of Earnings Mobility," *Econometrics Journal*, 11, 478-498.
- [35] Van Cleave, J., S.L. Gortmaker and J.M. Perrin (2010), "Dynamics of Obesity and Chronic Health Conditions Among Children and Youth," *Journal of the American Medical Association*, 303 (7), 623-630.
- [36] Walpole, S.C., D. Prieto-Merino, P. Edwards, J. Cleland, G. Stevens, and I. Roberts (2012), "The Weight of Nations: An Estimation of Adult Human Biomass," *BMC Public Health*, 12, 439.
- [37] Whitaker, R.C., J.A. Wright, M.S. Pepe, K.D. Seidel, and W.H. Dietz (1997), "Predicting Obesity in Young Adulthood from Childhood and Parental Obesity," *New England Journal of Medicine*, 337, 869-873.

Appendix A. Simulation Details.

- Case I. Own y_{i0} , own x_{it} , replace $\alpha_i = \bar{\alpha}$, set $\varepsilon_{it} = 0$. This eliminates heterogeneity due to time invariant and time-varying unobserved factors.

- Compute

$$\hat{y}_{it} = \hat{\gamma}\hat{y}_{it-1} + x_{it}\hat{\beta} + \bar{\alpha}, \quad t = 1, \dots, T$$

where $\hat{y}_{i0} = y_{i0}$.

- Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0) = \frac{\sum_i \mathbf{I}(\hat{y}_{iT} > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_i \mathbf{I}(y_{i0} \geq y_0)}.$$

- Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0, i \in G) = \frac{\sum_{i \in G} \mathbf{I}(\hat{y}_{iT} > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_{i \in G} \mathbf{I}(y_{i0} \geq y_0)},$$

where G denotes a specific sub-sample of the data (e.g., males).

- Case II. Own y_{i0} , own x_{it} , draw $\alpha_i \sim F(\alpha)$ where $F(\cdot)$ is the empirical distribution of α_i , set $\varepsilon_{it} = 0$. This allows for time invariant unobserved heterogeneity, but breaks the correlation between x and α .

- Draw $\tilde{\alpha}_i(r) \sim F(\hat{\alpha})$, $r = 1, \dots, R$, where R is the number of simulations.

- Compute

$$\hat{y}_{it}(r) = \hat{\gamma}\hat{y}_{it-1}(r) + x_{it}\hat{\beta} + \tilde{\alpha}_i(r), \quad t = 1, \dots, T$$

where $\hat{y}_{i0}(r) = y_{i0} \forall r$.

- Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0) = \frac{1}{R} \sum_r \left[\frac{\sum_i \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_i \mathbf{I}(y_{i0} \geq y_0)} \right].$$

- Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0, i \in G) = \frac{1}{R} \sum_r \left[\frac{\sum_{i \in G} \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_{i \in G} \mathbf{I}(y_{i0} \geq y_0)} \right].$$

- Case III. Own y_{i0} , own x_{it} , draw $\alpha_i \sim F_G(\alpha)$ where $F_G(\cdot)$ is the empirical distribution of α_i in sub-sample G and $i \in G$, set $\varepsilon_{it} = 0$. This only partially breaks the correlation between x and α as it retains any correlation between x and α common to group G .

- Draw $\tilde{\alpha}_i(r) \sim F_G(\hat{\alpha})$, where $i \in G$, $r = 1, \dots, R$, and R is the number of simulations.

- Compute

$$\hat{y}_{it}(r) = \hat{\gamma}\hat{y}_{it-1}(r) + x_{it}\hat{\beta} + \tilde{\alpha}_i, \quad t = 1, \dots, T$$

where $\hat{y}_{i0}(r) = y_{i0} \forall r$.

- Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0) = \frac{1}{R} \sum_r \left[\frac{\sum_i \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_i \mathbf{I}(y_{i0} \geq y_0)} \right].$$

(d) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0, i \in G) = \frac{1}{R} \sum_r \left[\frac{\sum_{i \in G} \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_{i \in G} \mathbf{I}(y_{i0} \geq y_0)} \right].$$

4. Case IV. Own y_{i0} , own x_{it} , draw $\alpha_i \sim F_{G'}(\alpha)$ where $F_{G'}(\cdot)$ is the empirical distribution of α_i in sub-sample G' and $i \notin G'$, set $\varepsilon_{it} = 0$. This replaces the true α_i for those in group G (e.g., males) with a draw from the empirical distribution in group G' (e.g., females).

(a) Draw $\tilde{\alpha}_i \sim F_{G'}(\hat{\alpha})$, where $i \notin G'$, $r = 1, \dots, R$, and R is the number of simulations.

(b) Compute

$$\hat{y}_{it} = \hat{\gamma} \hat{y}_{it-1} + x_{it} \hat{\beta} + \tilde{\alpha}_i, \quad t = 1, \dots, T$$

where $\hat{y}_{i0} = y_{i0}$.

(c) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0) = \frac{1}{R} \sum_r \left[\frac{\sum_i \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_i \mathbf{I}(y_{i0} \geq y_0)} \right].$$

(d) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0, i \in G) = \frac{1}{R} \sum_r \left[\frac{\sum_{i \in G} \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_{i \in G} \mathbf{I}(y_{i0} \geq y_0)} \right].$$

5. Case V. Own y_{i0} , own x_{it} , own w_i , set $\eta_i = 0$, set $\varepsilon_{it} = 0$. This eliminates heterogeneity due to time invariant and time-varying unobserved factors.

(a) Compute

$$\hat{y}_{it}(r) = \hat{\gamma} \hat{y}_{it-1} + x_{it} \hat{\beta} + w_i \hat{\delta}, \quad t = 1, \dots, T$$

where $\hat{y}_{i0} = y_{i0}$.

(b) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0) = \frac{\sum_i \mathbf{I}(\hat{y}_{iT} > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_i \mathbf{I}(y_{i0} \geq y_0)}.$$

(c) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0, i \in G) = \frac{\sum_{i \in G} \mathbf{I}(\hat{y}_{iT} > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_{i \in G} \mathbf{I}(y_{i0} \geq y_0)}.$$

6. Case VI. Own y_{i0} , own x_{it} , draw $w_i \sim F(w)$ where $F(\cdot)$ is the empirical distribution of w_i , set $\eta_i = 0$, set $\varepsilon_{it} = 0$. This eliminates heterogeneity due to time invariant and time-varying unobserved factors and breaks the correlation between x and w .

(a) Draw $\tilde{w}_i(r) \sim F(w)$, $r = 1, \dots, R$, where R is the number of simulations.

(b) Compute

$$\hat{y}_{it}(r) = \hat{\gamma} \hat{y}_{it-1}(r) + x_{it} \hat{\beta} + \tilde{w}_i(r) \hat{\delta}, \quad t = 1, \dots, T$$

where $\hat{y}_{i0}(r) = y_{i0} \forall r$.

(c) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0) = \frac{1}{R} \sum_r \left[\frac{\sum_i \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_i \mathbf{I}(y_{i0} \geq y_0)} \right].$$

(d) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0, i \in G) = \frac{1}{R} \sum_r \left[\frac{\sum_{i \in G} \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_{i \in G} \mathbf{I}(y_{i0} \geq y_0)} \right].$$

7. Case VII. Own y_{i0} , own x_{it} , draw $w_i \sim F_G(w)$ where $F_G(\cdot)$ is the empirical distribution of w_i in sub-sample G and $i \in G$, set $\eta_i = 0$, set $\varepsilon_{it} = 0$. This eliminates heterogeneity due to time invariant and time-varying unobserved factors and only partially breaks the correlation between x and α as it retains any correlation between x and w common to group G .

(a) Draw $\tilde{w}_i(r) \sim F_G(w)$, where $i \in G$, $r = 1, \dots, R$, and R is the number of simulations.

(b) Compute

$$\hat{y}_{it}(r) = \hat{\gamma} \hat{y}_{it-1}(r) + x_{it} \hat{\beta} + \tilde{w}_i(r) \hat{\delta}, \quad t = 1, \dots, T$$

where $\hat{y}_{i0}(r) = y_{i0} \forall r$.

(c) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0) = \frac{1}{R} \sum_r \left[\frac{\sum_i \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_i \mathbf{I}(y_{i0} \geq y_0)} \right].$$

(d) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0, i \in G) = \frac{1}{R} \sum_r \left[\frac{\sum_{i \in G} \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_{i \in G} \mathbf{I}(y_{i0} \geq y_0)} \right].$$

8. Case VIII. Own y_{i0} , own x_{it} , draw $w_i \sim F_{G'}(w)$ where $F_{G'}(\cdot)$ is the empirical distribution of w_i in sub-sample G' and $i \notin G$, set $\eta_i = 0$, set $\varepsilon_{it} = 0$. This replaces the true w_i for those in group G (e.g., males) with a draw from the empirical distribution in group G' (e.g., females) and eliminates heterogeneity due to time invariant and time-varying unobserved factors.

(a) Draw $\tilde{w}_i(r) \sim F_{G'}(w)$, where $i \notin G$, $r = 1, \dots, R$, and R is the number of simulations.

(b) Compute

$$\hat{y}_{it}(r) = \hat{\gamma} \hat{y}_{it-1}(r) + x_{it} \hat{\beta} + \tilde{w}_i(r) \hat{\delta}, \quad t = 1, \dots, T$$

where $\hat{y}_{i0}(r) = y_{i0} \forall r$.

(c) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0) = \frac{1}{R} \sum_r \left[\frac{\sum_i \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_i \mathbf{I}(y_{i0} > y_0)} \right].$$

(d) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0, i \in G) = \frac{1}{R} \sum_r \left[\frac{\sum_{i \in G} \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_{i \in G} \mathbf{I}(y_{i0} > y_0)} \right].$$

9. Case IX. Own y_{i0} , own x_{it} , own w_i , draw $\eta_i \sim F(\eta)$ where $F(\cdot)$ is the empirical distribution of η_i , set $\varepsilon_{it} = 0$. This breaks the correlation between x , w and η and eliminates heterogeneity due to time-varying unobserved factors.

(a) Draw $\tilde{\eta}_i(r) \sim F(\hat{\eta})$, $r = 1, \dots, R$, where R is the number of simulations.

(b) Compute

$$\hat{y}_{it}(r) = \hat{\gamma}\hat{y}_{it-1}(r) + x_{it}\hat{\beta} + w_i\hat{\delta} + \tilde{\eta}_i(r), \quad t = 1, \dots, T$$

where $\hat{y}_{i0}(r) = y_{i0} \forall r$.

(c) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0) = \frac{1}{R} \sum_r \left[\frac{\sum_i \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_i \mathbf{I}(y_{i0} \geq y_0)} \right].$$

(d) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0, i \in G) = \frac{1}{R} \sum_r \left[\frac{\sum_{i \in G} \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_{i \in G} \mathbf{I}(y_{i0} \geq y_0)} \right].$$

10. Case X. Own y_{i0} , own x_{it} , own w_i , draw $\eta_i \sim F_G(\eta)$ where $F_G(\cdot)$ is the empirical distribution of η_i in sub-sample G and $i \in G$, set $\varepsilon_{it} = 0$. This only partially breaks the correlation between x , w and η as it retains any correlation between x , w and η common to group G and eliminates heterogeneity due to time invariant and time-varying unobserved factors.

(a) Draw $\tilde{\eta}_i(r) \sim F_G(\hat{\eta})$, where $i \in G$, $r = 1, \dots, R$, and R is the number of simulations.

(b) Compute

$$\hat{y}_{it}(r) = \hat{\gamma}\hat{y}_{it-1}(r) + x_{it}\hat{\beta} + w_i\hat{\delta} + \tilde{\eta}_i(r), \quad t = 1, \dots, T$$

where $\hat{y}_{i0}(r) = y_{i0} \forall r$.

(c) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0) = \frac{1}{R} \sum_r \left[\frac{\sum_i \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_i \mathbf{I}(y_{i0} \geq y_0)} \right].$$

(d) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0, i \in G) = \frac{1}{R} \sum_r \left[\frac{\sum_{i \in G} \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_{i \in G} \mathbf{I}(y_{i0} \geq y_0)} \right].$$

11. Case XI. Own y_{i0} , own x_{it} , own w_i , draw $\eta_i \sim F_{G'}(\eta)$ where $F_{G'}(\cdot)$ is the empirical distribution of η_i in sub-sample G' and $i \notin G$, set $\varepsilon_{it} = 0$. This replaces the true η_i for those in group G (e.g., males) with a draw from the empirical distribution in group G' (e.g., females) and eliminates heterogeneity due to time invariant and time-varying unobserved factors.

(a) Draw $\tilde{\eta}_i(r) \sim F_{G'}(\hat{\eta})$, where $i \notin G$, $r = 1, \dots, R$, and R is the number of simulations.

(b) Compute

$$\hat{y}_{it}(r) = \hat{\gamma}\hat{y}_{it-1}(r) + x_{it}\hat{\beta} + w_i\hat{\delta} + \tilde{\eta}_i(r), \quad t = 1, \dots, T$$

where $\hat{y}_{i0}(r) = y_{i0} \forall r$.

(c) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0) = \frac{1}{R} \sum_r \left[\frac{\sum_i \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_i \mathbf{I}(y_{i0} > y_0)} \right].$$

(d) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0, i \in G) = \frac{1}{R} \sum_r \left[\frac{\sum_{i \in G} \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_{i \in G} \mathbf{I}(y_{i0} > y_0)} \right].$$

12. Case XII. Own y_{i0} , replace $x_{it} = \bar{x}_t$, own α_i , set $\varepsilon_{it} = 0$. This eliminates heterogeneity due to time-varying factors.

(a) Compute

$$\hat{y}_{it} = \hat{\gamma}\hat{y}_{it-1} + \bar{x}_t\hat{\beta} + \hat{\alpha}_i, \quad t = 1, \dots, T$$

where $\hat{y}_{i0} = y_{i0}$.

(b) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0) = \frac{\sum_i \mathbf{I}(\hat{y}_{iT} > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_i \mathbf{I}(y_{i0} \geq y_0)}.$$

(c) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0, i \in G) = \frac{\sum_{i \in G} \mathbf{I}(\hat{y}_{iT} > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_{i \in G} \mathbf{I}(y_{i0} \geq y_0)},$$

where G denotes a specific sub-sample of the data (e.g., males).

13. Case XIII. Own y_{i0} , draw $x_i \sim F(x_1, \dots, x_T)$ where $F(\cdot)$ is the empirical joint distribution of x_1, \dots, x_T , own α_i , set $\varepsilon_{it} = 0$. This breaks the correlation between x and α and eliminates heterogeneity due to time-varying unobserved factors.

(a) Draw $\tilde{x}_i(r) \sim F(x_1, \dots, x_T)$, $r = 1, \dots, R$, where R is the number of simulations.

(b) Compute

$$\hat{y}_{it}(r) = \hat{\gamma}\hat{y}_{it-1}(r) + \tilde{x}_{it}(r)\hat{\beta} + \alpha_i, \quad t = 1, \dots, T$$

where $\hat{y}_{i0}(r) = y_{i0} \forall r$.

(c) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0) = \frac{1}{R} \sum_r \left[\frac{\sum_i \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_i \mathbf{I}(y_{i0} \geq y_0)} \right].$$

(d) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0, i \in G) = \frac{1}{R} \sum_r \left[\frac{\sum_{i \in G} \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_{i \in G} \mathbf{I}(y_{i0} \geq y_0)} \right].$$

14. Case XIV. Own y_{i0} , draw $x_i \sim F_G(x_1, \dots, x_T)$ where $F_G(\cdot)$ is the empirical joint distribution of x_1, \dots, x_T in sub-sample G and $i \in G$, own α_i , set $\varepsilon_{it} = 0$. This only partially breaks the correlation between x and α as it retains any correlation between x and α common to group G and eliminates heterogeneity due to time-varying unobserved factors.

(a) Draw $\tilde{x}_i(r) \sim F_G(x_1, \dots, x_T)$, where $i \in G$, $r = 1, \dots, R$, where R is the number of simulations.

(b) Compute

$$\hat{y}_{it}(r) = \hat{\gamma}\hat{y}_{it-1}(r) + \tilde{x}_{it}(r)\hat{\beta} + \alpha_i, \quad t = 1, \dots, T$$

where $\hat{y}_{i0}(r) = y_{i0} \forall r$.

(c) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0) = \frac{1}{R} \sum_r \left[\frac{\sum_i \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_i \mathbf{I}(y_{i0} > y_0)} \right].$$

(d) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0, i \in G) = \frac{1}{R} \sum_r \left[\frac{\sum_{i \in G} \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_{i \in G} \mathbf{I}(y_{i0} > y_0)} \right].$$

15. Case XV. Own y_{i0} , draw $x_i. \sim F_G(x_1, \dots, x_T)$ where $F_G(\cdot)$ is the empirical joint distribution of x_1, \dots, x_T in sub-sample G and $i \in G$, own α_i , set $\varepsilon_{it} = 0$. This replaces the true $x_i.$ for those in group G (e.g., males) with a draw from the empirical distribution in group G' (e.g., females) and eliminates heterogeneity due to time-varying unobserved factors.

(a) Draw $\tilde{x}_i.(r) \sim F_{G'}(x_1, \dots, x_T)$, where $i \notin G'$, $r = 1, \dots, R$, where R is the number of simulations.

(b) Compute

$$\hat{y}_{it}(r) = \hat{\gamma} \hat{y}_{it-1}(r) + \tilde{x}_{it}(r) \hat{\beta} + \alpha_i, \quad t = 1, \dots, T$$

where $\hat{y}_{i0}(r) = y_{i0} \forall r$.

(c) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0) = \frac{1}{R} \sum_r \left[\frac{\sum_i \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_i \mathbf{I}(y_{i0} > y_0)} \right].$$

(d) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0, i \in G) = \frac{1}{R} \sum_r \left[\frac{\sum_{i \in G} \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_{i \in G} \mathbf{I}(y_{i0} > y_0)} \right].$$

16. Case XVI. Own y_{i0} , own x_{it} , own α_i , draw $\varepsilon_i. \sim F(\varepsilon_1, \dots, \varepsilon_T)$ where $F(\cdot)$ is the empirical distribution of $\varepsilon_i.$. This breaks the correlation between α and ε .

(a) Draw $\tilde{\varepsilon}_i.(r) \sim F(\hat{\varepsilon}_1, \dots, \hat{\varepsilon}_T)$, $r = 1, \dots, R$, where R is the number of simulations.

(b) Compute

$$\hat{y}_{it}(r) = \hat{\gamma} \hat{y}_{it-1}(r) + x_{it} \hat{\beta} + \hat{\alpha}_i + \tilde{\varepsilon}_{it}(r), \quad t = 1, \dots, T$$

where $\hat{y}_{i0}(r) = y_{i0} \forall r$.

(c) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0) = \frac{1}{R} \sum_r \left[\frac{\sum_i \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} \geq y_0)}{\sum_i \mathbf{I}(y_{i0} \geq y_0)} \right].$$

(d) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0, i \in G) = \frac{1}{R} \sum_r \left[\frac{\sum_{i \in G} \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} \geq y_0)}{\sum_{i \in G} \mathbf{I}(y_{i0} \geq y_0)} \right].$$

17. Case XVII. Own y_{i0} , own x_{it} , own α_i , draw $\varepsilon_i. \sim F_G(\varepsilon_1, \dots, \varepsilon_T)$ where $F_G(\cdot)$ is the empirical distribution of $\varepsilon_i.$ in sub-sample G and $i \in G$. This only partially breaks the correlation between α and ε as it retains any correlation between α and ε common to group G .

(a) Draw $\tilde{\varepsilon}_i.(r) \sim F_G(\hat{\varepsilon}_1, \dots, \hat{\varepsilon}_T)$, where $i \in G$, $r = 1, \dots, R$, and R is the number of simulations.

(b) Compute

$$\hat{y}_{it}(r) = \hat{\gamma} \hat{y}_{it-1}(r) + x_{it} \hat{\beta} + \hat{\alpha}_i + \tilde{\varepsilon}_{it}(r), \quad t = 1, \dots, T$$

where $\hat{y}_{i0}(r) = y_{i0} \forall r$.

(c) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0) = \frac{1}{R} \sum_r \left[\frac{\sum_i \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_i \mathbf{I}(y_{i0} > y_0)} \right].$$

(d) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0, i \in G) = \frac{1}{R} \sum_r \left[\frac{\sum_{i \in G} \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_{i \in G} \mathbf{I}(y_{i0} > y_0)} \right].$$

18. Case XVIII. Own y_{i0} , own x_{it} , own α_i , draw $\varepsilon_i \sim F_{G'}(\varepsilon_1, \dots, \varepsilon_T)$ where $F_{G'}(\cdot)$ is the empirical distribution of ε_i in sub-sample G' and $i \notin G$. This replaces the true ε_i for those in group G (e.g., males) with a draw from the empirical distribution in group G' (e.g., females).

(a) Draw $\tilde{\varepsilon}_i(r) \sim F_{G'}(\hat{\varepsilon}_1, \dots, \hat{\varepsilon}_T)$, where $i \notin G$, $r = 1, \dots, R$, and R is the number of simulations.

(b) Compute

$$\hat{y}_{it}(r) = \hat{\gamma} \hat{y}_{it-1}(r) + x_{it} \hat{\beta} + \hat{\alpha}_i + \tilde{\varepsilon}_{it}(r), \quad t = 1, \dots, T$$

where $\hat{y}_{i0}(r) = y_{i0} \forall r$.

(c) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0) = \frac{1}{R} \sum_r \left[\frac{\sum_i \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} \geq y_0)}{\sum_i \mathbf{I}(y_{i0} \geq y_0)} \right].$$

(d) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0, i \in G) = \frac{1}{R} \sum_r \left[\frac{\sum_{i \in G} \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} \geq y_0)}{\sum_{i \in G} \mathbf{I}(y_{i0} \geq y_0)} \right].$$

19. Case XIX. Own y_{i0} , own α_i , draw $x_i, \varepsilon_i \sim F(x_1, \dots, x_T, \varepsilon_1, \dots, \varepsilon_T)$ where $F(\cdot)$ is the empirical joint distribution of $x_1, \dots, x_T, \varepsilon_1, \dots, \varepsilon_T$. This breaks the correlation between α and x, ε .

(a) Draw $\tilde{x}_i(r), \tilde{\varepsilon}_i(r) \sim F(x_1, \dots, x_T, \hat{\varepsilon}_1, \dots, \hat{\varepsilon}_T)$, $r = 1, \dots, R$, where R is the number of simulations.

(b) Compute

$$\hat{y}_{it}(r) = \hat{\gamma} \hat{y}_{it-1}(r) + \tilde{x}_{it}(r) \hat{\beta} + \hat{\alpha}_i + \tilde{\varepsilon}_{it}(r), \quad t = 1, \dots, T$$

where $\hat{y}_{i0}(r) = y_{i0} \forall r$.

(c) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0) = \frac{1}{R} \sum_r \left[\frac{\sum_i \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} \geq y_0)}{\sum_i \mathbf{I}(y_{i0} \geq y_0)} \right].$$

(d) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0, i \in G) = \frac{1}{R} \sum_r \left[\frac{\sum_{i \in G} \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} \geq y_0)}{\sum_{i \in G} \mathbf{I}(y_{i0} \geq y_0)} \right].$$

20. Case XX. Own y_{i0} , own α_i , draw $x_i, \varepsilon_i \sim F_G(x_1, \dots, x_T, \varepsilon_1, \dots, \varepsilon_T)$ where $F_G(\cdot)$ is the empirical joint distribution of $x_1, \dots, x_T, \varepsilon_1, \dots, \varepsilon_T$ in sub-sample G and $i \in G$. This only partially breaks the correlation between α and x, ε as it retains any correlation between α and x, ε common to group G .

(a) Draw $\tilde{x}_i(r), \tilde{\varepsilon}_i(r) \sim F_G(x_1, \dots, x_T, \hat{\varepsilon}_1, \dots, \hat{\varepsilon}_T)$, where $i \in G$, $r = 1, \dots, R$, and R is the number of simulations.

(b) Compute

$$\hat{y}_{it}(r) = \hat{\gamma} \hat{y}_{it-1}(r) + \tilde{x}_{it}(r) \hat{\beta} + \hat{\alpha}_i + \tilde{\varepsilon}_{it}(r), \quad t = 1, \dots, T$$

where $\hat{y}_{i0}(r) = y_{i0} \forall r$.

(c) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0) = \frac{1}{R} \sum_r \left[\frac{\sum_i \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_i \mathbf{I}(y_{i0} > y_0)} \right].$$

(d) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0, i \in G) = \frac{1}{R} \sum_r \left[\frac{\sum_{i \in G} \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} > y_0)}{\sum_{i \in G} \mathbf{I}(y_{i0} > y_0)} \right].$$

21. Case XXI. Own y_{i0} , own α_i , draw $x_i, \varepsilon_i \sim F(x_1, \dots, x_T, \varepsilon_1, \dots, \varepsilon_T)$ where $F_{G'}(\cdot)$ is the empirical joint distribution of $x_1, \dots, x_T, \varepsilon_1, \dots, \varepsilon_T$ in sub-sample G' and $i \notin G$. This replaces the true x_i and ε_i for those in group G (e.g., males) with a draw from the empirical distribution in group G' (e.g., females).

(a) Draw $\tilde{x}_i(r), \tilde{\varepsilon}_i(r) \sim F_{G'}(x_1, \dots, x_T, \hat{\varepsilon}_1, \dots, \hat{\varepsilon}_T)$, where $i \notin G, r = 1, \dots, R$, and R is the number of simulations.

(b) Compute

$$\hat{y}_{it}(r) = \hat{\gamma} \hat{y}_{it-1}(r) + \tilde{x}_{it}(r) \hat{\beta} + \hat{\alpha}_i + \tilde{\varepsilon}_{it}(r), \quad t = 1, \dots, T$$

where $\hat{y}_{i0}(r) = y_{i0} \forall r$.

(c) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0) = \frac{1}{R} \sum_r \left[\frac{\sum_i \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} \geq y_0)}{\sum_i \mathbf{I}(y_{i0} \geq y_0)} \right].$$

(d) Compute

$$\Pr(y_{iT} > y^* | y_{i0} \geq y_0, i \in G) = \frac{1}{R} \sum_r \left[\frac{\sum_{i \in G} \mathbf{I}(\hat{y}_{iT}(r) > y^*) \mathbf{I}(y_{i0} \geq y_0)}{\sum_{i \in G} \mathbf{I}(y_{i0} \geq y_0)} \right].$$

Appendix B. Figures.




Figure B1. Weight: Decomposition.

Notes: Data from ECLS-K. Empirical distributions of the outcome, y_{it} , and the various estimated components of the dynamic model $y_{it} = \gamma y_{it-1} + X_{it}\beta + \alpha_i + \varepsilon_{it}$. Data from ECLS-K. See text for further details.




Figure B2. Weight: Decomposition of Fixed Effects.

Notes: Data from ECLS-K. Empirical distributions of the estimated fixed effects, α_{it} , and its components obtained from the dynamic model $y_{it} = \gamma y_{it-1} + X_{it}\beta + \alpha_i + \varepsilon_{it}$, where $\alpha_i = w_i\delta + \eta_i$. See text for further details.




Figure B3. Height: Decomposition.

Notes: See Figure B1 and text for details.




Figure B4. Height: Decomposition of Fixed Effects.

Notes: See Figure B2 and text for details.




Figure B5. BMI: Decomposition.

Notes: See Figure B1 and text for details.




Figure B6. BMI: Decomposition of Fixed Effects.

Notes: See Figure B2 and text for details.




Figure B7. Weight: Decomposition.

Notes: Data from ECLS-B. Empirical distributions of the outcome, y_{it} , and the various estimated components of the dynamic model $y_{it} = \gamma y_{it-1} + X_{it}\beta + \alpha_i + \varepsilon_{it}$. See text for further details.




Figure B8. Weight: Decomposition of Fixed Effects.

Notes: Data from ECLS-B. Empirical distributions of the estimated fixed effects, α_{it} , and its components obtained from the dynamic model $y_{it} = \gamma y_{it-1} + X_{it}\beta + \alpha_i + \varepsilon_{it}$, where $\alpha_i = w_i\delta + \eta_i$. See text for further details.




Figure B9. Height: Decomposition.

Notes: See Figure B7 and text for details.




Figure B10. Height: Decomposition of Fixed Effects.
 Notes: See Figure B8 and text for details.

Table 1. Dynamic Panel Data Estimates: Weight Z-Scores.

	Full Sample			Race						Gender					
				White		Non-White				Male			Female		
Lag Weight	0.931*	0.932*	0.775*	0.932*	0.932*	0.686*	0.929*	0.931*	0.895*	0.948*	0.951*	0.276*	0.914*	0.910*	1.732*
	(0.003)	(0.003)	(0.067)	(0.004)	(0.004)	(0.080)	(0.004)	(0.004)	(0.116)	(0.004)	(0.004)	(0.056)	(0.004)	(0.004)	(0.219)
Time-Varying Covariates	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes
Time Invariant Covariates	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No
Fixed Effects	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes
Observations	27470	27470	27470	16900	16900	16900	10570	10570	10570	13880	13880	13880	13580	13580	13580
Underidentification	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000
Endogeneity	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.186	p = 0.050	p = 0.000	p = 0.113	p = 0.007	p = 0.000	p = 0.000	p = 0.000	p = 0.000
First-Stage F-stat	88146.9	84006.9	269.0	49635.2	48281.7	172.4	37871.6	35837.8	101.4	37300.4	36297.2	195.2	55116.6	51193.2	80.0

	Urban Status						Mother's Education						SES Status					
	Urban			Non-Urban			Less Than College			College			Low SES			High SES		
Lag Weight	0.929*	0.930*	0.896*	0.933*	0.932*	0.708*	0.931*	0.932*	0.743*	0.930*	0.930*	0.887*	0.929*	0.934*	0.807*	0.931*	0.931*	0.778*
	(0.004)	(0.005)	(0.116)	(0.003)	(0.004)	(0.082)	(0.003)	(0.003)	(0.077)	(0.006)	(0.006)	(0.146)	(0.005)	(0.005)	(0.135)	(0.003)	(0.003)	(0.077)
Time-Varying Covariates	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes
Time Invariant Covariates	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No
Fixed Effects	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes
Observations	10010	10010	10010	17460	17460	17460	20250	20250	20250	7210	7210	7210	8340	8340	8340	19120	19120	19120
Underidentification	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000
Endogeneity	p = 0.060	p = 0.006	p = 0.000	p = 0.003	p = 0.000	p = 0.000	p = 0.002	p = 0.000	p = 0.000	p = 0.005	p = 0.001	p = 0.000	p = 0.102	p = 0.036	p = 0.000	p = 0.000	p = 0.000	p = 0.000
First-Stage F-stat	33299.8	31697.0	103.1	54867.6	52628.1	167.7	65410.2	62000.1	186.1	23916.2	22706.4	78.0	28392.3	26903.2	65.3	59580.3	57058.5	212.0

Notes: ‡ p<0.10, † p<0.05, * p<0.01. Robust standard errors in parentheses. Estimation by GMM. Excluded instrument is the dependent variable twice-lagged. Sample sizes rounded to the nearest 10 per NCES restricted data regulations. Sample includes data from fall kindergarten, spring first, spring third, spring fifth grades, and spring eighth grade. See text for the list of covariates and further details.

Table 2. Dynamic Simulations: Weight Z-Scores, $\Pr(y_{it} \geq 85^{\text{th}} \text{ percentile} \mid y_{it} \geq 85^{\text{th}} \text{ percentile})$.

	Full	Race		Gender		Urban Status		Education		SES	
	Sample	White	Non-White	Male	Female	Urban	Non-Urban	Non-College	College	Low SES	High SES
Benchmark	0.840	0.823	0.861	0.846	0.833	0.840	0.840	0.870	0.748	0.880	0.820
Panel I. Own Xs, $\varepsilon=0$, and											
$\alpha = E[\alpha]$	0.753	0.681	0.846	0.765	0.739	0.776	0.740	0.800	0.609	0.868	0.695
$\alpha \sim f(\alpha)$	0.576	0.555	0.604	0.581	0.570	0.586	0.570	0.588	0.540	0.609	0.559
$\alpha \sim f_i(\alpha)$		0.552	0.608	0.597	0.555	0.565	0.584	0.607	0.484	0.637	0.547
$\alpha \sim f_{-i}(\alpha)$		0.562	0.602	0.567	0.586	0.599	0.551	0.537	0.561	0.597	0.593
Panel II. Own Xs, $\eta=0$, $\varepsilon=0$, and											
$W = W_i$	0.727	0.642	0.837	0.800	0.645	0.714	0.734	0.829	0.418	0.894	0.643
$W \sim f(W)$	0.703	0.649	0.775	0.710	0.696	0.729	0.689	0.736	0.604	0.786	0.662
$W \sim f_i(W)$		0.632	0.789	0.778	0.622	0.672	0.724	0.802	0.394	0.870	0.614
$W \sim f_{-i}(W)$		0.676	0.765	0.640	0.767	0.759	0.631	0.550	0.678	0.748	0.769
Panel III. Own Xs, own Ws, $\varepsilon=0$, and											
$\eta \sim f(\eta)$	0.591	0.563	0.628	0.616	0.563	0.582	0.596	0.622	0.497	0.652	0.561
$\eta \sim f_i(\eta)$		0.566	0.620	0.608	0.569	0.580	0.596	0.617	0.500	0.642	0.562
$\eta \sim f_{-i}(\eta)$		0.558	0.632	0.625	0.557	0.582	0.593	0.635	0.496	0.653	0.557
Panel IV. Own α, $\varepsilon=0$, and											
$X = E[X]$	0.844	0.834	0.858	0.850	0.838	0.837	0.848	0.867	0.777	0.867	0.833
$X \sim f(X)$	0.849	0.841	0.860	0.855	0.843	0.841	0.854	0.871	0.783	0.870	0.839
$X \sim f_i(X)$		0.834	0.868	0.854	0.844	0.844	0.852	0.873	0.771	0.877	0.834
$X \sim f_{-i}(X)$		0.852	0.856	0.856	0.843	0.839	0.857	0.864	0.788	0.866	0.852
Panel V. Own Xs, own α, and											
$\varepsilon \sim f(\varepsilon)$	0.846	0.830	0.868	0.854	0.838	0.842	0.849	0.873	0.766	0.880	0.830
$\varepsilon \sim f_i(\varepsilon)$		0.831	0.866	0.852	0.841	0.844	0.848	0.873	0.766	0.881	0.829
$\varepsilon \sim f_{-i}(\varepsilon)$		0.829	0.868	0.857	0.834	0.840	0.852	0.873	0.765	0.880	0.832
Panel VI. Own α and											
$X, \varepsilon \sim f(X, \varepsilon)$	0.843	0.835	0.854	0.851	0.835	0.833	0.849	0.866	0.775	0.863	0.833
$X, \varepsilon \sim f_i(X, \varepsilon)$		0.829	0.862	0.848	0.840	0.839	0.846	0.869	0.761	0.873	0.827
$X, \varepsilon \sim f_{-i}(X, \varepsilon)$		0.845	0.849	0.854	0.831	0.830	0.855	0.859	0.779	0.860	0.848

Notes: Benchmark case denotes the observed probability in the sample. Simulations - based on 500 draws - are obtained after estimating the dynamic model $y_{it} = \alpha_i + \gamma y_{it-1} + X_{it}\beta + \varepsilon_{it}$, where $\alpha_i = w_i\delta + \eta_i$. $f(\cdot)$ denotes the empirical distribution of the argument. Sample includes data from fall kindergarten, spring first, spring third, spring fifth grades, and spring eighth grade. See text for the list of covariates and further details.

Table 3. Dynamic Simulations: Weight Z-Scores, $\Pr(y_{it} \geq 95^{\text{th}} \text{ percentile} \mid y_{it} \geq 95^{\text{th}} \text{ percentile})$.

	Full	Race		Gender		Urban Status		Education		SES	
	Sample	White	Non-White	Male	Female	Urban	Non-Urban	Non-College	College	Low SES	High SES
Benchmark	0.762	0.732	0.795	0.807	0.710	0.791	0.746	0.790	0.646	0.799	0.740
Panel I. Own Xs, $\varepsilon=0$, and											
$\alpha=E[\alpha]$	0.134	0.093	0.180	0.171	0.091	0.150	0.125	0.151	0.067	0.183	0.105
$\alpha \sim f(\alpha)$	0.396	0.376	0.418	0.404	0.387	0.408	0.390	0.370	0.403	0.420	0.382
$\alpha \sim f_i(\alpha)$		0.368	0.431	0.434	0.356	0.386	0.403	0.300	0.428	0.461	0.364
$\alpha \sim f_{-i}(\alpha)$		0.390	0.411	0.377	0.418	0.420	0.371	0.398	0.337	0.405	0.425
Panel II. Own Xs, $\eta=0$, $\varepsilon=0$, and											
$W=W_i$	0.246	0.194	0.302	0.305	0.175	0.196	0.273	0.079	0.286	0.348	0.185
$W \sim f(W)$	0.196	0.152	0.245	0.221	0.167	0.218	0.185	0.135	0.212	0.253	0.163
$W \sim f_i(W)$		0.140	0.268	0.265	0.121	0.168	0.209	0.057	0.248	0.330	0.138
$W \sim f_{-i}(W)$		0.172	0.231	0.177	0.211	0.244	0.142	0.161	0.109	0.222	0.220
Panel III. Own Xs, own Ws, $\varepsilon=0$, and											
$\eta \sim f(\eta)$	0.418	0.397	0.441	0.443	0.388	0.406	0.425	0.332	0.439	0.464	0.391
$\eta \sim f_i(\eta)$		0.395	0.443	0.444	0.384	0.408	0.426	0.318	0.441	0.467	0.388
$\eta \sim f_{-i}(\eta)$		0.405	0.439	0.442	0.397	0.404	0.427	0.341	0.435	0.465	0.398
Panel IV. Own α, $\varepsilon=0$, and											
$X=E[X]$	0.780	0.774	0.786	0.809	0.746	0.791	0.773	0.691	0.801	0.796	0.770
$X \sim f(X)$	0.784	0.777	0.792	0.811	0.753	0.792	0.780	0.695	0.806	0.800	0.775
$X \sim f_i(X)$		0.768	0.806	0.811	0.753	0.797	0.778	0.682	0.810	0.816	0.769
$X \sim f_{-i}(X)$		0.792	0.784	0.813	0.751	0.788	0.786	0.700	0.795	0.793	0.791
Panel V. Own Xs, own α, and											
$\varepsilon \sim f(\varepsilon)$	0.786	0.763	0.811	0.813	0.754	0.806	0.775	0.678	0.812	0.820	0.766
$\varepsilon \sim f_i(\varepsilon)$		0.765	0.809	0.811	0.759	0.811	0.773	0.679	0.812	0.822	0.766
$\varepsilon \sim f_{-i}(\varepsilon)$		0.762	0.812	0.817	0.749	0.806	0.778	0.679	0.812	0.820	0.769
Panel VI. Own α and											
$X, \varepsilon \sim f(X, \varepsilon)$	0.779	0.769	0.791	0.806	0.747	0.785	0.776	0.685	0.802	0.797	0.768
$X, \varepsilon \sim f_i(X, \varepsilon)$		0.762	0.804	0.803	0.755	0.794	0.772	0.669	0.808	0.816	0.760
$X, \varepsilon \sim f_{-i}(X, \varepsilon)$		0.784	0.785	0.811	0.742	0.780	0.786	0.691	0.792	0.791	0.789

Notes: See Table 2 and text for further details.

Table 4. Dynamic Simulations: Weight Z-Scores, $\Pr(y_{it} \geq 85^{\text{th}} \text{ percentile} \mid y_{it} \leq 50^{\text{th}} \text{ percentile})$.

	Full	Race		Gender		Urban Status		Education		SES	
	Sample	White	Non-White	Male	Female	Urban	Non-Urban	Non-College	College	Low SES	High SES
Benchmark	0.118	0.121	0.113	0.132	0.104	0.108	0.124	0.131	0.079	0.145	0.106
Panel I. Own Xs, $\varepsilon=0$, and											
$\alpha=E[\alpha]$	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
$\alpha \sim f(\alpha)$	0.213	0.209	0.220	0.213	0.214	0.218	0.210	0.209	0.215	0.224	0.208
$\alpha \sim f_i(\alpha)$		0.198	0.238	0.244	0.182	0.205	0.218	0.146	0.238	0.264	0.192
$\alpha \sim f_{-i}(\alpha)$		0.226	0.210	0.181	0.245	0.226	0.198	0.232	0.151	0.206	0.247
Panel II. Own Xs, $\eta=0$, $\varepsilon=0$, and											
$W=W_i$	0.006	0.005	0.009	0.007	0.005	0.002	0.009	0.001	0.008	0.010	0.005
$W \sim f(W)$	0.006	0.006	0.008	0.006	0.006	0.007	0.006	0.005	0.007	0.008	0.006
$W \sim f_i(W)$		0.005	0.008	0.008	0.005	0.002	0.009	0.002	0.008	0.012	0.005
$W \sim f_{-i}(W)$		0.006	0.007	0.005	0.008	0.010	0.002	0.007	0.002	0.007	0.008
Panel III. Own Xs, own Ws, $\varepsilon=0$, and											
$\eta \sim f(\eta)$	0.209	0.204	0.216	0.225	0.193	0.198	0.215	0.166	0.223	0.240	0.195
$\eta \sim f_i(\eta)$		0.197	0.228	0.240	0.176	0.201	0.214	0.145	0.231	0.256	0.188
$\eta \sim f_{-i}(\eta)$		0.216	0.209	0.211	0.210	0.197	0.218	0.173	0.203	0.235	0.211
Panel IV. Own α, $\varepsilon=0$, and											
$X=E[X]$	0.116	0.123	0.105	0.137	0.096	0.100	0.126	0.085	0.127	0.134	0.109
$X \sim f(X)$	0.120	0.127	0.109	0.140	0.101	0.104	0.130	0.087	0.132	0.140	0.112
$X \sim f_i(X)$		0.123	0.115	0.140	0.101	0.106	0.128	0.083	0.134	0.148	0.109
$X \sim f_{-i}(X)$		0.135	0.106	0.141	0.101	0.103	0.133	0.088	0.125	0.136	0.119
Panel V. Own Xs, own α, and											
$\varepsilon \sim f(\varepsilon)$	0.121	0.125	0.116	0.140	0.103	0.109	0.129	0.080	0.135	0.149	0.109
$\varepsilon \sim f_i(\varepsilon)$		0.126	0.115	0.139	0.105	0.110	0.128	0.079	0.136	0.151	0.109
$\varepsilon \sim f_{-i}(\varepsilon)$		0.124	0.116	0.143	0.101	0.108	0.131	0.080	0.135	0.149	0.111
Panel VI. Own α and											
$X, \varepsilon \sim f(X, \varepsilon)$	0.124	0.131	0.112	0.143	0.105	0.108	0.133	0.088	0.136	0.143	0.115
$X, \varepsilon \sim f_i(X, \varepsilon)$		0.127	0.117	0.140	0.107	0.111	0.130	0.082	0.138	0.153	0.111
$X, \varepsilon \sim f_{-i}(X, \varepsilon)$		0.138	0.109	0.145	0.103	0.106	0.138	0.090	0.129	0.139	0.125

Notes: See Table 2 and text for further details.

Table 5. Dynamic Panel Data Estimates: Height Z-Scores.

	Full Sample			Race						Gender					
				White			Non-White			Male			Female		
Lag Height	0.937*	0.936*	0.603*	0.938*	0.941*	0.553*	0.932*	0.932*	0.676*	0.951*	0.954*	0.460*	0.922*	0.918*	0.739*
	(0.004)	(0.004)	(0.048)	(0.004)	(0.004)	(0.057)	(0.007)	(0.007)	(0.082)	(0.004)	(0.005)	(0.055)	(0.006)	(0.006)	(0.079)
Time-Varying Covariates	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes
Time Invariant Covariates	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No
Fixed Effects	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes
Observations	27470	27470	27470	16900	16900	16900	10570	10570	10570	13880	13880	13880	13580	13580	13580
Underidentification	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000
Endogeneity	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000
First-Stage F-stat	68631.4	64548.2	673.1	42314.9	40485.2	388.5	26601.1	24081.3	284.7	34323.0	31415.4	326.6	34269.4	32561.6	339.9

	Urban Status						Mother's Education						SES Status					
	Urban			Non-Urban			Less Than College			College			Low SES			High SES		
Lag Height	0.923*	0.922*	0.646*	0.945*	0.945*	0.580*	0.936*	0.936*	0.659*	0.939*	0.939*	0.453*	0.932*	0.933*	0.705*	0.938*	0.939*	0.560*
	(0.006)	(0.007)	(0.073)	(0.005)	(0.005)	(0.064)	(0.004)	(0.004)	(0.059)	(0.007)	(0.007)	(0.081)	(0.007)	(0.008)	(0.096)	(0.004)	(0.004)	(0.054)
Time-Varying Covariates	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes
Time Invariant Covariates	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No
Fixed Effects	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes
Observations	10010	10010	10010	17460	17460	17460	20250	20250	20250	7210	7210	7210	8340	8340	8340	19120	19120	19120
Underidentification	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000
Endogeneity	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000
First-Stage F-stat	20304.1	19094.2	270.5	50866.0	47176.7	405.4	54613.3	51233.9	498.4	14521.0	13396.4	166.2	21762.4	20541.3	211.9	46682.7	44002.2	464.8

Notes: ‡ p<0.10, † p<0.05, * p<0.01. See Table 1 and text for further details.

Table 6. Dynamic Simulations: Height Z-Scores, $\Pr(y_{iT} \geq 85^{\text{th}} \text{ percentile} \mid y_{i1} \geq 85^{\text{th}} \text{ percentile})$.

	Full	Race		Gender		Urban Status		Education		SES	
	Sample	White	Non-White	Male	Female	Urban	Non-Urban	Non-College	College	Low SES	High SES
Benchmark	0.606	0.635	0.559	0.665	0.545	0.592	0.614	0.587	0.653	0.515	0.642
Panel I. Own Xs, $\varepsilon=0$, and											
$\alpha=E[\alpha]$	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
$\alpha \sim f(\alpha)$	0.235	0.243	0.221	0.236	0.234	0.229	0.238	0.228	0.252	0.222	0.240
$\alpha \sim f_i(\alpha)$		0.248	0.216	0.260	0.209	0.221	0.243	0.228	0.251	0.217	0.243
$\alpha \sim f_{-i}(\alpha)$		0.237	0.223	0.210	0.259	0.236	0.228	0.228	0.250	0.222	0.234
Panel II. Own Xs, $\eta=0$, $\varepsilon=0$, and											
$W=W_i$	0.006	0.006	0.008	0.010	0.002	0.008	0.005	0.004	0.012	0.002	0.008
$W \sim f(W)$	0.003	0.004	0.002	0.004	0.003	0.004	0.003	0.002	0.005	0.002	0.004
$W \sim f_i(W)$		0.003	0.004	0.006	0.002	0.003	0.003	0.002	0.007	0.002	0.004
$W \sim f_{-i}(W)$		0.007	0.001	0.002	0.004	0.005	0.002	0.003	0.006	0.002	0.004
Panel III. Own Xs, own Ws, $\varepsilon=0$, and											
$\eta \sim f(\eta)$	0.256	0.265	0.241	0.281	0.230	0.246	0.261	0.247	0.279	0.231	0.266
$\eta \sim f_i(\eta)$		0.263	0.241	0.286	0.224	0.245	0.261	0.250	0.270	0.237	0.263
$\eta \sim f_{-i}(\eta)$		0.267	0.240	0.276	0.234	0.245	0.260	0.234	0.283	0.225	0.271
Panel IV. Own α, $\varepsilon=0$, and											
$X=E[X]$	0.669	0.662	0.680	0.702	0.635	0.673	0.667	0.667	0.673	0.627	0.686
$X \sim f(X)$	0.663	0.655	0.675	0.695	0.629	0.664	0.662	0.664	0.660	0.626	0.677
$X \sim f_i(X)$		0.669	0.655	0.694	0.630	0.657	0.665	0.655	0.686	0.611	0.686
$X \sim f_{-i}(X)$		0.633	0.688	0.697	0.627	0.668	0.656	0.688	0.650	0.634	0.657
Panel V. Own Xs, own α, and											
$\varepsilon \sim f(\varepsilon)$	0.635	0.645	0.621	0.665	0.604	0.627	0.640	0.627	0.656	0.581	0.657
$\varepsilon \sim f_i(\varepsilon)$		0.667	0.587	0.680	0.590	0.616	0.645	0.622	0.670	0.562	0.667
$\varepsilon \sim f_{-i}(\varepsilon)$		0.610	0.641	0.650	0.620	0.631	0.632	0.644	0.648	0.590	0.635
Panel VI. Own α and											
$X, \varepsilon \sim f(X, \varepsilon)$	0.635	0.629	0.646	0.666	0.603	0.636	0.635	0.638	0.630	0.603	0.648
$X, \varepsilon \sim f_i(X, \varepsilon)$		0.663	0.595	0.680	0.589	0.621	0.644	0.624	0.668	0.569	0.666
$X, \varepsilon \sim f_{-i}(X, \varepsilon)$		0.572	0.679	0.653	0.616	0.643	0.620	0.675	0.616	0.618	0.608

Notes: Benchmark case denotes the observed probability in the sample. Simulations - **based on 500 draws** - obtained after estimating the dynamic model $y_{it} = \alpha_i + \gamma y_{it-1} + X_{it}\beta + \varepsilon_{it}$, where $\alpha_i = w_i\delta + \eta_i$. $f(\cdot)$ denotes the empirical distribution of the argument. Sample includes data from fall kindergarten, spring first, spring third, spring fifth grades, and spring eighth grade. See text for the list of covariates and further details.

Table 7. Dynamic Simulations: Height Z-Scores, $\Pr(y_{iT} \geq 95^{\text{th}} \text{ percentile} \mid y_{i1} \geq 95^{\text{th}} \text{ percentile})$.

	Full	Race		Gender		Urban Status		Education		SES	
	Sample	White	Non-White	Male	Female	Urban	Non-Urban	Non-College	College	Low SES	High SES
Benchmark	0.467	0.481	0.446	0.550	0.377	0.448	0.476	0.453	0.502	0.414	0.489
Panel I. Own Xs, $\varepsilon=0$, and											
$\alpha=E[\alpha]$	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
$\alpha \sim f(\alpha)$	0.092	0.097	0.085	0.094	0.090	0.092	0.092	0.103	0.088	0.083	0.096
$\alpha \sim f_i(\alpha)$		0.097	0.086	0.107	0.076	0.087	0.095	0.103	0.087	0.082	0.097
$\alpha \sim f_{-i}(\alpha)$		0.096	0.081	0.079	0.102	0.095	0.087	0.104	0.087	0.084	0.091
Panel II. Own Xs, $\eta=0$, $\varepsilon=0$, and											
$W=W_i$	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
$W \sim f(W)$	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
$W \sim f_i(W)$		0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
$W \sim f_{-i}(W)$		0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Panel III. Own Xs, own Ws, $\varepsilon=0$, and											
$\eta \sim f(\eta)$	0.108	0.111	0.103	0.122	0.092	0.107	0.108	0.121	0.102	0.094	0.113
$\eta \sim f_i(\eta)$		0.110	0.105	0.127	0.090	0.108	0.108	0.112	0.105	0.098	0.111
$\eta \sim f_{-i}(\eta)$		0.116	0.099	0.120	0.093	0.107	0.107	0.122	0.095	0.093	0.118
Panel IV. Own α, $\varepsilon=0$, and											
$X=E[X]$	0.523	0.498	0.562	0.572	0.469	0.493	0.536	0.498	0.533	0.517	0.525
$X \sim f(X)$	0.523	0.495	0.568	0.569	0.473	0.499	0.534	0.500	0.532	0.520	0.524
$X \sim f_i(X)$		0.509	0.540	0.566	0.475	0.492	0.539	0.530	0.522	0.497	0.533
$X \sim f_{-i}(X)$		0.470	0.585	0.570	0.472	0.503	0.526	0.488	0.562	0.531	0.502
Panel V. Own Xs, own α, and											
$\varepsilon \sim f(\varepsilon)$	0.506	0.496	0.522	0.545	0.463	0.480	0.518	0.505	0.506	0.485	0.514
$\varepsilon \sim f_i(\varepsilon)$		0.517	0.487	0.559	0.449	0.470	0.523	0.523	0.499	0.466	0.524
$\varepsilon \sim f_{-i}(\varepsilon)$		0.461	0.543	0.530	0.477	0.484	0.508	0.498	0.521	0.496	0.493
Panel VI. Own α and											
$X, \varepsilon \sim f(X, \varepsilon)$	0.509	0.481	0.552	0.546	0.468	0.489	0.518	0.481	0.519	0.509	0.509
$X, \varepsilon \sim f_i(X, \varepsilon)$		0.517	0.494	0.558	0.457	0.476	0.528	0.523	0.506	0.470	0.528
$X, \varepsilon \sim f_{-i}(X, \varepsilon)$		0.425	0.587	0.533	0.479	0.497	0.502	0.467	0.561	0.525	0.468

Notes: See Table 6 and text for further details.

Table 8. Dynamic Simulations: Height Z-Scores, $\Pr(y_{iT} \geq 85^{\text{th}} \text{ percentile} \mid y_{i1} \leq 50^{\text{th}} \text{ percentile})$.

	Full	Race		Gender		Urban Status		Education		SES	
	Sample	White	Non-White	Male	Female	Urban	Non-Urban	Non-College	College	Low SES	High SES
Benchmark	0.030	0.036	0.019	0.032	0.027	0.025	0.032	0.030	0.029	0.026	0.032
Panel I. Own Xs, $\varepsilon=0$, and											
$\alpha=E[\alpha]$	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
$\alpha \sim f(\alpha)$	0.144	0.151	0.134	0.144	0.145	0.141	0.146	0.158	0.140	0.136	0.149
$\alpha \sim f_i(\alpha)$		0.153	0.135	0.163	0.126	0.134	0.151	0.157	0.141	0.134	0.150
$\alpha \sim f_{-i}(\alpha)$		0.151	0.135	0.125	0.164	0.145	0.140	0.160	0.138	0.135	0.147
Panel II. Own Xs, $\eta=0$, $\varepsilon=0$, and											
$W=W_i$	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
$W \sim f(W)$	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
$W \sim f_i(W)$		0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
$W \sim f_{-i}(W)$		0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Panel III. Own Xs, own Ws, $\varepsilon=0$, and											
$\eta \sim f(\eta)$	0.133	0.145	0.115	0.149	0.117	0.125	0.138	0.155	0.126	0.114	0.142
$\eta \sim f_i(\eta)$		0.143	0.119	0.153	0.115	0.126	0.138	0.144	0.130	0.119	0.139
$\eta \sim f_{-i}(\eta)$		0.149	0.114	0.145	0.120	0.125	0.139	0.159	0.118	0.112	0.149
Panel IV. Own α, $\varepsilon=0$, and											
$X=E[X]$	0.019	0.020	0.017	0.019	0.019	0.021	0.018	0.014	0.020	0.020	0.018
$X \sim f(X)$	0.018	0.019	0.016	0.017	0.018	0.019	0.017	0.014	0.019	0.019	0.017
$X \sim f_i(X)$		0.020	0.014	0.017	0.018	0.018	0.017	0.015	0.018	0.018	0.018
$X \sim f_{-i}(X)$		0.017	0.017	0.017	0.018	0.019	0.017	0.013	0.021	0.020	0.015
Panel V. Own Xs, own α, and											
$\varepsilon \sim f(\varepsilon)$	0.021	0.024	0.016	0.022	0.020	0.021	0.021	0.021	0.021	0.021	0.021
$\varepsilon \sim f_i(\varepsilon)$		0.026	0.014	0.022	0.020	0.020	0.021	0.023	0.020	0.019	0.022
$\varepsilon \sim f_{-i}(\varepsilon)$		0.021	0.018	0.021	0.020	0.021	0.020	0.021	0.023	0.021	0.019
Panel VI. Own α and											
$X, \varepsilon \sim f(X, \varepsilon)$	0.022	0.024	0.019	0.023	0.021	0.023	0.022	0.019	0.023	0.023	0.022
$X, \varepsilon \sim f_i(X, \varepsilon)$		0.027	0.015	0.023	0.021	0.022	0.022	0.024	0.021	0.020	0.023
$X, \varepsilon \sim f_{-i}(X, \varepsilon)$		0.019	0.022	0.023	0.021	0.023	0.021	0.018	0.028	0.024	0.018

Notes: See Table 6 and text for further details.

Table 9. Dynamic Panel Data Estimates: Body Mass Index Z-Scores.

	Full Sample			Race						Gender					
				White			Non-White			Male			Female		
Lag BMI	0.912*	0.911*	0.217*	0.915*	0.912*	0.194*	0.904*	0.910*	0.255*	0.915*	0.919*	0.179*	0.909*	0.903*	0.275*
	(0.004)	(0.005)	(0.015)	(0.006)	(0.006)	(0.018)	(0.007)	(0.007)	(0.027)	(0.007)	(0.007)	(0.018)	(0.006)	(0.006)	(0.026)
Time-Varying Covariates	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes
Time Invariant Covariates	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No
Fixed Effects	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes
Observations	27470	27470	27470	16900	16900	16900	10570	10570	10570	13880	13880	13880	13580	13580	13580
Underidentification	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000
Endogeneity	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000
First-Stage F-stat	8637.8	8518.0	820.8	4744.4	4885.9	535.8	3823.7	3657.3	293.8	3555.1	3516.1	451.8	6391.0	6229.0	420.2

	Urban Status						Mother's Education						SES Status					
	Urban			Non-Urban			Less Than College			College			Low SES			High SES		
Lag BMI	0.912*	0.909*	0.254*	0.912*	0.911*	0.200*	0.906*	0.908*	0.216*	0.921*	0.920*	0.227*	0.905*	0.912*	0.222*	0.912*	0.910*	0.222*
	(0.007)	(0.007)	(0.027)	(0.006)	(0.006)	(0.018)	(0.005)	(0.005)	(0.017)	(0.009)	(0.009)	(0.030)	(0.008)	(0.008)	(0.028)	(0.006)	(0.006)	(0.018)
Time-Varying Covariates	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes
Time Invariant Covariates	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No
Fixed Effects	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes
Observations	10010	10010	10010	17460	17460	17460	20250	20250	20250	7210	7210	7210	8340	8340	8340	19120	19120	19120
Underidentification	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000
Endogeneity	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000
First-Stage F-stat	3452.7	3187.6	278.1	5265.0	5388.1	545.0	6257.6	6090.8	575.5	2448.2	2562.1	254.0	3340.8	3264.5	251.2	5367.1	5346.3	579.1

Notes: ‡ p<0.10, † p<0.05, * p<0.01. See Table 1 and text for further details.

Table 10. Dynamic Simulations: BMI Z-Scores, $\Pr(y_{it} \geq 85^{\text{th}} \text{ percentile} \mid y_{it} \geq 85^{\text{th}} \text{ percentile})$.

	Full	Race		Gender		Urban Status		Education		SES	
	Sample	White	Non-White	Male	Female	Urban	Non-Urban	Non-College	College	Low SES	High SES
Benchmark	0.746	0.703	0.800	0.736	0.757	0.758	0.739	0.779	0.637	0.813	0.710
Panel I. Own Xs, $\varepsilon=0$, and											
$\alpha=E[\alpha]$	0.000	0.001	0.000	0.000	0.001	0.000	0.001	0.000	0.002	0.000	0.001
$\alpha \sim f(\alpha)$	0.347	0.348	0.345	0.346	0.347	0.347	0.347	0.357	0.344	0.341	0.350
$\alpha \sim f_i(\alpha)$		0.309	0.407	0.357	0.337	0.344	0.348	0.265	0.376	0.420	0.316
$\alpha \sim f_{-i}(\alpha)$		0.410	0.307	0.334	0.360	0.348	0.345	0.392	0.250	0.306	0.430
Panel II. Own Xs, $\eta=0$, $\varepsilon=0$, and											
$W=W_i$	0.070	0.011	0.145	0.076	0.065	0.079	0.066	0.004	0.090	0.147	0.030
$W \sim f(W)$	0.055	0.056	0.055	0.054	0.057	0.054	0.056	0.071	0.051	0.049	0.059
$W \sim f_i(W)$		0.015	0.118	0.065	0.044	0.059	0.052	0.006	0.067	0.108	0.028
$W \sim f_{-i}(W)$		0.121	0.015	0.041	0.069	0.050	0.062	0.093	0.004	0.023	0.130
Panel III. Own Xs, own Ws, $\varepsilon=0$, and											
$\eta \sim f(\eta)$	0.368	0.333	0.412	0.376	0.359	0.370	0.366	0.299	0.389	0.413	0.344
$\eta \sim f_i(\eta)$		0.324	0.421	0.379	0.356	0.373	0.365	0.277	0.394	0.433	0.333
$\eta \sim f_{-i}(\eta)$		0.347	0.406	0.374	0.363	0.370	0.369	0.306	0.378	0.406	0.364
Panel IV. Own α, $\varepsilon=0$, and											
$X=E[X]$	0.797	0.761	0.842	0.793	0.802	0.798	0.797	0.708	0.824	0.845	0.772
$X \sim f(X)$	0.795	0.759	0.840	0.791	0.800	0.798	0.793	0.707	0.822	0.844	0.769
$X \sim f_i(X)$		0.761	0.837	0.789	0.802	0.798	0.793	0.723	0.818	0.837	0.773
$X \sim f_{-i}(X)$		0.756	0.841	0.792	0.798	0.798	0.793	0.701	0.831	0.847	0.760
Panel V. Own Xs, own α, and											
$\varepsilon \sim f(\varepsilon)$	0.766	0.733	0.807	0.764	0.768	0.768	0.764	0.688	0.789	0.807	0.744
$\varepsilon \sim f_i(\varepsilon)$		0.733	0.806	0.752	0.780	0.770	0.762	0.670	0.793	0.816	0.739
$\varepsilon \sim f_{-i}(\varepsilon)$		0.732	0.807	0.775	0.754	0.767	0.767	0.696	0.775	0.802	0.755
Panel VI. Own α and											
$X, \varepsilon \sim f(X, \varepsilon)$	0.765	0.730	0.808	0.763	0.766	0.768	0.762	0.672	0.792	0.813	0.739
$X, \varepsilon \sim f_i(X, \varepsilon)$		0.732	0.806	0.750	0.781	0.771	0.761	0.670	0.793	0.817	0.737
$X, \varepsilon \sim f_{-i}(X, \varepsilon)$		0.726	0.811	0.777	0.752	0.768	0.764	0.674	0.789	0.812	0.741

Notes: Benchmark case denotes the observed probability in the sample. Simulations - based on 500 draws - obtained after estimating the dynamic model $y_{it} = \alpha_i + \gamma y_{it-1} + X_{it}\beta + \varepsilon_{it}$, where $\alpha_i = w_i\delta + \eta_i$. $f(\cdot)$ denotes the empirical distribution of the argument. Sample includes data from fall kindergarten, spring first, spring third, spring fifth grades, and spring eighth grade. See text for the list of covariates and further details.

Table 11. Dynamic Simulations: BMI Z-Scores, $\Pr(y_{it} \geq 95^{\text{th}} \text{ percentile} \mid y_{it} \geq 95^{\text{th}} \text{ percentile})$.

	Full	Race		Gender		Urban Status		Education		SES	
	Sample	White	Non-White	Male	Female	Urban	Non-Urban	Non-College	College	Low SES	High SES
Benchmark	0.715	0.664	0.769	0.724	0.703	0.738	0.702	0.757	0.538	0.783	0.672
Panel I. Own Xs, $\varepsilon=0$, and											
$\alpha=E[\alpha]$	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
$\alpha \sim f(\alpha)$	0.179	0.180	0.178	0.178	0.180	0.177	0.179	0.187	0.177	0.175	0.181
$\alpha \sim f_i(\alpha)$		0.147	0.228	0.194	0.161	0.181	0.176	0.117	0.199	0.233	0.153
$\alpha \sim f_{-i}(\alpha)$		0.231	0.144	0.158	0.196	0.176	0.179	0.211	0.109	0.149	0.242
Panel II. Own Xs, $\eta=0$, $\varepsilon=0$, and											
$W=W_i$	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
$W \sim f(W)$	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
$W \sim f_i(W)$		0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
$W \sim f_{-i}(W)$		0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Panel III. Own Xs, own Ws, $\varepsilon=0$, and											
$\eta \sim f(\eta)$	0.195	0.171	0.221	0.199	0.190	0.195	0.196	0.137	0.209	0.226	0.176
$\eta \sim f_i(\eta)$		0.162	0.238	0.207	0.182	0.198	0.193	0.124	0.216	0.244	0.168
$\eta \sim f_{-i}(\eta)$		0.184	0.212	0.192	0.197	0.194	0.198	0.142	0.190	0.217	0.193
Panel IV. Own α, $\varepsilon=0$, and											
$X=E[X]$	0.792	0.757	0.830	0.804	0.778	0.810	0.782	0.629	0.832	0.845	0.759
$X \sim f(X)$	0.792	0.758	0.827	0.802	0.778	0.808	0.782	0.628	0.831	0.841	0.761
$X \sim f_i(X)$		0.760	0.824	0.801	0.781	0.808	0.783	0.648	0.826	0.830	0.766
$X \sim f_{-i}(X)$		0.754	0.830	0.805	0.776	0.809	0.782	0.622	0.845	0.846	0.749
Panel V. Own Xs, own α, and											
$\varepsilon \sim f(\varepsilon)$	0.734	0.699	0.771	0.739	0.728	0.748	0.726	0.597	0.767	0.776	0.708
$\varepsilon \sim f_i(\varepsilon)$		0.698	0.770	0.723	0.746	0.752	0.724	0.575	0.774	0.788	0.700
$\varepsilon \sim f_{-i}(\varepsilon)$		0.698	0.771	0.755	0.710	0.747	0.729	0.606	0.747	0.770	0.721
Panel VI. Own α and											
$X, \varepsilon \sim f(X, \varepsilon)$	0.735	0.699	0.773	0.742	0.726	0.749	0.727	0.583	0.771	0.784	0.704
$X, \varepsilon \sim f_i(X, \varepsilon)$		0.700	0.770	0.724	0.743	0.750	0.727	0.579	0.772	0.787	0.702
$X, \varepsilon \sim f_{-i}(X, \varepsilon)$		0.694	0.775	0.761	0.706	0.749	0.729	0.583	0.767	0.783	0.706

Notes: See Table 10 and text for further details.

Table 12. Dynamic Simulations: BMI Z-Scores, $\Pr(y_{it} \geq 85^{\text{th}} \text{ percentile} \mid y_{it} \leq 50^{\text{th}} \text{ percentile})$.

	Full	Race		Gender		Urban Status		Education		SES	
	White	White	Non-White	Male	Female	Urban	Non-Urban	Non-College	College	Low SES	High SES
Benchmark	0.142	0.127	0.167	0.152	0.132	0.138	0.144	0.162	0.087	0.192	0.121
Panel I. Own Xs, $\varepsilon=0$, and											
$\alpha=E[\alpha]$	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
$\alpha \sim f(\alpha)$	0.346	0.347	0.343	0.345	0.347	0.346	0.346	0.356	0.342	0.339	0.348
$\alpha \sim f_i(\alpha)$		0.308	0.403	0.354	0.335	0.344	0.347	0.263	0.375	0.418	0.315
$\alpha \sim f_{-i}(\alpha)$		0.410	0.304	0.333	0.359	0.348	0.342	0.390	0.249	0.304	0.428
Panel II. Own Xs, $\eta=0$, $\varepsilon=0$, and											
$W=W_i$	0.044	0.020	0.086	0.051	0.036	0.032	0.050	0.005	0.058	0.094	0.023
$W \sim f(W)$	0.054	0.054	0.053	0.052	0.055	0.054	0.053	0.066	0.049	0.047	0.056
$W \sim f_i(W)$		0.014	0.114	0.063	0.042	0.061	0.049	0.005	0.065	0.105	0.026
$W \sim f_{-i}(W)$		0.118	0.014	0.040	0.067	0.051	0.058	0.087	0.003	0.021	0.126
Panel III. Own Xs, own Ws, $\varepsilon=0$, and											
$\eta \sim f(\eta)$	0.335	0.311	0.378	0.345	0.326	0.328	0.339	0.270	0.360	0.392	0.312
$\eta \sim f_i(\eta)$		0.300	0.387	0.347	0.320	0.329	0.337	0.248	0.363	0.410	0.303
$\eta \sim f_{-i}(\eta)$		0.324	0.370	0.341	0.328	0.325	0.341	0.276	0.345	0.383	0.332
Panel IV. Own α, $\varepsilon=0$, and											
$X=E[X]$	0.105	0.096	0.120	0.120	0.091	0.092	0.113	0.062	0.121	0.139	0.091
$X \sim f(X)$	0.107	0.098	0.121	0.121	0.092	0.095	0.113	0.062	0.123	0.142	0.092
$X \sim f_i(X)$		0.099	0.121	0.121	0.093	0.095	0.113	0.067	0.121	0.138	0.094
$X \sim f_{-i}(X)$		0.097	0.122	0.122	0.092	0.095	0.113	0.060	0.129	0.144	0.089
Panel V. Own Xs, own α, and											
$\varepsilon \sim f(\varepsilon)$	0.125	0.116	0.139	0.138	0.111	0.115	0.130	0.086	0.139	0.158	0.111
$\varepsilon \sim f_i(\varepsilon)$		0.116	0.139	0.131	0.117	0.116	0.129	0.079	0.142	0.163	0.109
$\varepsilon \sim f_{-i}(\varepsilon)$		0.116	0.138	0.145	0.105	0.114	0.131	0.088	0.131	0.155	0.115
Panel VI. Own α and											
$X, \varepsilon \sim f(X, \varepsilon)$	0.125	0.116	0.140	0.140	0.111	0.115	0.131	0.083	0.141	0.161	0.111
$X, \varepsilon \sim f_i(X, \varepsilon)$		0.116	0.140	0.132	0.117	0.116	0.130	0.081	0.141	0.162	0.110
$X, \varepsilon \sim f_{-i}(X, \varepsilon)$		0.116	0.141	0.147	0.105	0.114	0.132	0.083	0.139	0.161	0.111

Notes: See Table 10 and text for further details.

Table 13. Dynamic Panel Data Estimates: Weight Z-Scores.

	Full Sample			Race						Gender					
				White			Non-White			Male			Female		
Lag Weight	0.873*	0.870*	0.124*	0.868*	0.903*	0.105*	0.873*	0.857*	0.144*	0.888*	0.896*	0.108*	0.857*	0.850*	0.143*
	(0.010)	(0.012)	(0.013)	(0.016)	(0.021)	(0.018)	(0.013)	(0.016)	(0.018)	(0.014)	(0.019)	(0.019)	(0.014)	(0.017)	(0.018)
Time-Varying Covariates	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes
Time Invariant Covariates	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No
Fixed Effects	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes
Observations	10900	10900	10900	4500	4500	4500	6400	6400	6400	5450	5450	5450	5400	5400	5400
Underidentification	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000
Endogeneity	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000
First-Stage F-stat	112.0	1398.4	1929.3	626.3	429.3	803.7	1683.9	1112.7	1253.0	321.8	778.8	972.4	889.5	640.4	965.5

	Urban Status						Mother's Education						SES Status					
	Urban			Non-Urban			Less Than College			College			Low SES			High SES		
Lag Weight	0.874*	0.869*	0.130*	0.870*	0.894*	0.121*	0.864*	0.886*	0.123*	0.875*	0.872*	0.126*	0.868*	0.887*	0.127*	0.874*	0.860*	0.128*
	(0.012)	(0.015)	(0.016)	(0.018)	(0.024)	(0.023)	(0.018)	(0.024)	(0.023)	(0.012)	(0.015)	(0.016)	(0.018)	(0.021)	(0.022)	(0.012)	(0.015)	(0.016)
Time-Varying Covariates	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes
Time Invariant Covariates	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No
Fixed Effects	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes
Observations	7800	7800	7800	3100	3100	3100	7750	7750	7750	3150	3150	3150	4050	4050	4050	6850	6850	6850
Underidentification	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000
Endogeneity	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000
First-Stage F-stat	1785.2	1184.3	1566.7	469.1	342.0	497.2	697.1	447.3	715.3	1455.9	978.3	1261.1	686.6	481.3	687.5	1497.8	988.9	1339.1

Notes: ‡ p<0.10, † p<0.05, * p<0.01. Robust standard errors in parentheses. Estimation by GMM. Excluded instrument is the dependent variable twice-lagged. Sample sizes rounded to the nearest 50 per NCES restricted data regulations. Sample includes data from waves 1-4 in the ECLS-B. See text for the list of covariates and further details.

Table 14. Dynamic Simulations: Weight Z-Scores, $\Pr(y_{it} \geq 85^{\text{th}} \text{ percentile} \mid y_{it} \geq 85^{\text{th}} \text{ percentile})$.

	Full	Race		Gender		Urban Status		Education		SES	
	Sample	White	Non-White	Male	Female	Urban	Non-Urban	Non-College	College	Low SES	High SES
Benchmark	0.541	0.508	0.558	0.605	0.496	0.511	0.621	0.434	0.583	0.593	0.507
Panel I. Own Xs, $\varepsilon=0$, and											
$\alpha=E[\alpha]$	0.007	0.000	0.010	0.005	0.008	0.006	0.008	0.000	0.009	0.014	0.002
$\alpha \sim f(\alpha)$	0.255	0.284	0.240	0.249	0.259	0.240	0.295	0.237	0.262	0.257	0.254
$\alpha \sim f_i(\alpha)$		0.216	0.284	0.263	0.243	0.248	0.267	0.213	0.271	0.285	0.237
$\alpha \sim f_{-i}(\alpha)$		0.333	0.180	0.234	0.275	0.217	0.308	0.246	0.242	0.240	0.284
Panel II. Own Xs, $\eta=0$, $\varepsilon=0$, and											
$W=W_i$	0.169	0.135	0.187	0.181	0.161	0.157	0.202	0.112	0.191	0.174	0.166
$W \sim f(W)$	0.101	0.120	0.092	0.090	0.109	0.085	0.142	0.080	0.109	0.106	0.098
$W \sim f_i(W)$		0.050	0.127	0.103	0.094	0.094	0.107	0.068	0.115	0.122	0.087
$W \sim f_{-i}(W)$		0.166	0.042	0.078	0.121	0.064	0.154	0.086	0.093	0.096	0.114
Panel III. Own Xs, own Ws, $\varepsilon=0$, and											
$\eta \sim f(\eta)$	0.332	0.311	0.343	0.340	0.326	0.322	0.358	0.304	0.343	0.338	0.328
$\eta \sim f_i(\eta)$		0.307	0.347	0.344	0.326	0.324	0.354	0.294	0.348	0.357	0.318
$\eta \sim f_{-i}(\eta)$		0.320	0.337	0.340	0.327	0.320	0.361	0.309	0.336	0.328	0.346
Panel IV. Own α, $\varepsilon=0$, and											
$X=E[X]$	0.605	0.532	0.642	0.678	0.554	0.593	0.637	0.558	0.624	0.646	0.578
$X \sim f(X)$	0.596	0.521	0.634	0.659	0.552	0.591	0.611	0.557	0.612	0.630	0.574
$X \sim f_i(X)$		0.560	0.609	0.656	0.557	0.576	0.652	0.525	0.623	0.639	0.568
$X \sim f_{-i}(X)$		0.494	0.670	0.662	0.550	0.630	0.593	0.570	0.585	0.628	0.582
Panel V. Own Xs, own α, and											
$\varepsilon \sim f(\varepsilon)$	0.581	0.556	0.594	0.635	0.543	0.556	0.649	0.516	0.607	0.611	0.562
$\varepsilon \sim f_i(\varepsilon)$		0.553	0.595	0.630	0.547	0.553	0.653	0.511	0.607	0.616	0.557
$\varepsilon \sim f_{-i}(\varepsilon)$		0.558	0.592	0.640	0.538	0.560	0.647	0.520	0.603	0.608	0.567
Panel VI. Own α and											
$X, \varepsilon \sim f(X, \varepsilon)$	0.582	0.509	0.618	0.633	0.546	0.578	0.593	0.546	0.596	0.615	0.561
$X, \varepsilon \sim f_i(X, \varepsilon)$		0.541	0.596	0.626	0.554	0.564	0.634	0.510	0.609	0.625	0.553
$X, \varepsilon \sim f_{-i}(X, \varepsilon)$		0.486	0.652	0.641	0.539	0.618	0.577	0.559	0.566	0.609	0.574

Notes: Benchmark case denotes the observed probability in the sample. Simulations - based on 500 draws - obtained after estimating the dynamic model $y_{it} = \alpha_i + \gamma y_{it-1} + X_{it}\beta + \varepsilon_{it}$, where $\alpha_i = w_i\delta + \eta_i$. $f(\cdot)$ denotes the empirical distribution of the argument. Sample includes data from waves 1-4 of the ECLS-B. See text for the list of covariates and further details.

Table 15. Dynamic Simulations: Weight Z-Scores, $\Pr(y_{it} \geq 95^{\text{th}} \text{ percentile} \mid y_{it} \geq 95^{\text{th}} \text{ percentile})$.

	Full	Race		Gender		Urban Status		Education		SES	
	Sample	White	Non-White	Male	Female	Urban	Non-Urban	Non-College	College	Low SES	High SES
Benchmark	0.366	0.360	0.369	0.409	0.340	0.354	0.392	0.198	0.420	0.446	0.302
Panel I. Own Xs, $\varepsilon=0$, and											
$\alpha=E[\alpha]$	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
$\alpha \sim f(\alpha)$	0.118	0.134	0.112	0.113	0.122	0.109	0.140	0.105	0.123	0.122	0.116
$\alpha \sim f_i(\alpha)$		0.088	0.139	0.121	0.112	0.114	0.132	0.084	0.132	0.150	0.101
$\alpha \sim f_{-i}(\alpha)$		0.166	0.070	0.105	0.131	0.098	0.147	0.112	0.102	0.104	0.145
Panel II. Own Xs, $\eta=0$, $\varepsilon=0$, and											
$W=W_i$	0.031	0.016	0.038	0.044	0.023	0.028	0.038	0.030	0.032	0.043	0.022
$W \sim f(W)$	0.007	0.007	0.007	0.006	0.007	0.006	0.009	0.006	0.007	0.008	0.005
$W \sim f_i(W)$		0.001	0.011	0.007	0.005	0.006	0.007	0.003	0.008	0.012	0.004
$W \sim f_{-i}(W)$		0.011	0.002	0.004	0.010	0.004	0.010	0.005	0.005	0.006	0.008
Panel III. Own Xs, own Ws, $\varepsilon=0$, and											
$\eta \sim f(\eta)$	0.166	0.148	0.174	0.174	0.162	0.162	0.177	0.150	0.172	0.171	0.163
$\eta \sim f_i(\eta)$		0.133	0.183	0.174	0.161	0.160	0.178	0.131	0.179	0.190	0.151
$\eta \sim f_{-i}(\eta)$		0.157	0.164	0.171	0.164	0.162	0.178	0.158	0.156	0.158	0.185
Panel IV. Own α, $\varepsilon=0$, and											
$X=E[X]$	0.419	0.392	0.430	0.465	0.390	0.427	0.400	0.337	0.445	0.462	0.384
$X \sim f(X)$	0.416	0.381	0.431	0.448	0.396	0.427	0.390	0.335	0.442	0.455	0.385
$X \sim f_i(X)$		0.412	0.410	0.445	0.401	0.415	0.437	0.309	0.452	0.462	0.381
$X \sim f_{-i}(X)$		0.354	0.464	0.451	0.392	0.461	0.374	0.347	0.417	0.450	0.393
Panel V. Own Xs, own α, and											
$\varepsilon \sim f(\varepsilon)$	0.405	0.415	0.401	0.443	0.382	0.390	0.441	0.284	0.444	0.462	0.360
$\varepsilon \sim f_i(\varepsilon)$		0.414	0.403	0.438	0.387	0.386	0.450	0.275	0.447	0.467	0.356
$\varepsilon \sim f_{-i}(\varepsilon)$		0.417	0.398	0.448	0.379	0.393	0.441	0.288	0.440	0.456	0.369
Panel VI. Own α and											
$X, \varepsilon \sim f(X, \varepsilon)$	0.413	0.372	0.430	0.443	0.394	0.425	0.385	0.337	0.437	0.450	0.382
$X, \varepsilon \sim f_i(X, \varepsilon)$		0.398	0.413	0.437	0.402	0.412	0.427	0.303	0.450	0.464	0.374
$X, \varepsilon \sim f_{-i}(X, \varepsilon)$		0.356	0.456	0.453	0.385	0.458	0.368	0.352	0.407	0.441	0.397

Notes: See Table 14 and text for further details.

Table 16. Dynamic Simulations: Weight Z-Scores, $\Pr(y_{it} \geq 85^{\text{th}} \text{ percentile} \mid y_{it} \leq 50^{\text{th}} \text{ percentile})$.

	Full	Race		Gender		Urban Status		Education		SES	
	Sample	White	Non-White	Male	Female	Urban	Non-Urban	Non-College	College	Low SES	High SES
Benchmark	0.431	0.426	0.435	0.443	0.418	0.426	0.443	0.395	0.446	0.468	0.410
Panel I. Own Xs, $\varepsilon=0$, and											
$\alpha=E[\alpha]$	0.837	0.982	0.724	0.830	0.846	0.793	0.949	0.798	0.854	0.820	0.847
$\alpha \sim f(\alpha)$	0.617	0.646	0.593	0.614	0.619	0.605	0.645	0.594	0.626	0.627	0.610
$\alpha \sim f_i(\alpha)$		0.594	0.632	0.624	0.610	0.614	0.622	0.592	0.627	0.635	0.606
$\alpha \sim f_{-i}(\alpha)$		0.683	0.537	0.603	0.629	0.579	0.653	0.594	0.626	0.622	0.619
Panel II. Own Xs, $\eta=0$, $\varepsilon=0$, and											
$W=W_i$	0.613	0.553	0.661	0.633	0.591	0.613	0.615	0.559	0.636	0.648	0.594
$W \sim f(W)$	0.719	0.777	0.674	0.715	0.725	0.698	0.771	0.685	0.734	0.732	0.712
$W \sim f_i(W)$		0.680	0.749	0.734	0.706	0.715	0.737	0.676	0.737	0.740	0.708
$W \sim f_{-i}(W)$		0.847	0.566	0.696	0.743	0.659	0.784	0.688	0.727	0.726	0.721
Panel III. Own Xs, own Ws, $\varepsilon=0$, and											
$\eta \sim f(\eta)$	0.556	0.525	0.581	0.568	0.542	0.556	0.555	0.526	0.568	0.574	0.546
$\eta \sim f_i(\eta)$		0.527	0.577	0.569	0.543	0.559	0.551	0.532	0.567	0.578	0.543
$\eta \sim f_{-i}(\eta)$		0.523	0.585	0.570	0.541	0.552	0.557	0.525	0.575	0.572	0.551
Panel IV. Own α, $\varepsilon=0$, and											
$X=E[X]$	0.435	0.398	0.465	0.462	0.405	0.447	0.407	0.444	0.432	0.432	0.437
$X \sim f(X)$	0.440	0.401	0.470	0.464	0.412	0.451	0.413	0.442	0.439	0.442	0.439
$X \sim f_i(X)$		0.436	0.448	0.460	0.416	0.436	0.452	0.413	0.450	0.451	0.434
$X \sim f_{-i}(X)$		0.377	0.503	0.467	0.410	0.488	0.397	0.455	0.411	0.436	0.447
Panel V. Own Xs, own α, and											
$\varepsilon \sim f(\varepsilon)$	0.431	0.428	0.434	0.453	0.407	0.429	0.436	0.402	0.444	0.450	0.421
$\varepsilon \sim f_i(\varepsilon)$		0.425	0.437	0.449	0.411	0.428	0.440	0.396	0.446	0.458	0.417
$\varepsilon \sim f_{-i}(\varepsilon)$		0.431	0.430	0.457	0.403	0.435	0.434	0.405	0.438	0.447	0.429
Panel VI. Own α and											
$X, \varepsilon \sim f(X, \varepsilon)$	0.441	0.405	0.469	0.464	0.415	0.451	0.417	0.442	0.441	0.445	0.439
$X, \varepsilon \sim f_i(X, \varepsilon)$		0.431	0.451	0.457	0.423	0.436	0.455	0.408	0.454	0.460	0.429
$X, \varepsilon \sim f_{-i}(X, \varepsilon)$		0.387	0.496	0.472	0.407	0.488	0.401	0.456	0.410	0.437	0.454

Notes: See Table 14 and text for further details.

Table 17. Dynamic Panel Data Estimates: Height.

	Full Sample			Race						Gender					
				White		Non-White				Male			Female		
Lag Height	0.480*	0.506*	-0.002	0.488*	0.522*	0.002	0.474*	0.493*	-0.005	0.485*	0.511*	-0.056*	0.474*	0.498*	-0.043*
	(0.004)	(0.010)	(0.007)	(0.006)	(0.016)	(0.012)	(0.005)	(0.013)	(0.009)	(0.006)	(0.014)	(0.008)	(0.006)	(0.015)	(0.008)
Time-Varying Covariates	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes
Time Invariant Covariates	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No
Fixed Effects	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes
Observations	10900	10900	10900	4500	4500	4500	6400	6400	6400	5450	5450	5450	5400	5400	5400
Underidentification	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000
Endogeneity	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000
First-Stage F-stat	770.2	6940.2	10250.2	17737.1	2263.8	3947.3	27111.7	4758.6	6311.6	568.9	3435.1	8396.2	21349.5	3328.9	8158.5

	Urban Status						Mother's Education						SES Status					
	Urban			Non-Urban			Less Than College			College			Low SES			High SES		
Lag Height	0.475*	0.491*	-0.049*	0.493*	0.549*	-0.057*	0.481*	0.492*	-0.051*	0.480*	0.515*	-0.049*	0.466*	0.486*	-0.055*	0.488*	0.524*	-0.045*
	(0.005)	(0.012)	(0.007)	(0.008)	(0.021)	(0.011)	(0.007)	(0.018)	(0.010)	(0.005)	(0.012)	(0.007)	(0.007)	(0.017)	(0.009)	(0.005)	(0.013)	(0.007)
Time-Varying Covariates	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes	No	Yes	Yes
Time Invariant Covariates	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No
Fixed Effects	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes	No	No	Yes
Observations	7800	7800	7800	3100	3100	3100	7750	7750	7750	3150	3150	3150	4050	4050	4050	6850	6850	6850
Underidentification	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000
Endogeneity	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.006	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000	p = 0.000
First-Stage F-stat	33778.2	5408.6	12465.7	310.8	1489.6	3939.6	13369.8	1660.6	4565.1	31408.3	5230.3	11710.0	246.8	2752.0	6982.2	28553.6	4057.3	9316.1

Notes: ‡ p<0.10, † p<0.05, * p<0.01. Dependent variable is length/height in centimeters. See Table 13 and text for further details.

Table 18. Dynamic Simulations: Height, $\Pr(y_{it} \geq 85^{\text{th}} \text{ percentile} \mid y_{it} \geq 85^{\text{th}} \text{ percentile})$.

	Full	Race		Gender		Urban Status		Education		SES	
	Sample	White	Non-White	Male	Female	Urban	Non-Urban	Non-College	College	Low SES	High SES
Benchmark	0.600	0.576	0.615	0.626	0.580	0.611	0.574	0.585	0.606	0.605	0.598
Panel I. Own Xs, $\varepsilon=0$, and											
$\alpha=E[\alpha]$	0.190	0.225	0.169	0.062	0.292	0.175	0.228	0.135	0.213	0.198	0.186
$\alpha \sim f(\alpha)$	0.380	0.400	0.368	0.332	0.418	0.373	0.397	0.355	0.390	0.387	0.376
$\alpha \sim f_i(\alpha)$		0.351	0.403	0.368	0.381	0.385	0.368	0.377	0.381	0.383	0.379
$\alpha \sim f_{-i}(\alpha)$		0.435	0.318	0.295	0.456	0.342	0.410	0.345	0.416	0.392	0.369
Panel II. Own Xs, $\eta=0$, $\varepsilon=0$, and											
$W=W_i$	0.396	0.367	0.413	0.386	0.403	0.401	0.383	0.386	0.400	0.373	0.409
$W \sim f(W)$	0.292	0.323	0.274	0.210	0.357	0.281	0.320	0.252	0.308	0.306	0.284
$W \sim f_i(W)$		0.256	0.316	0.258	0.300	0.301	0.266	0.275	0.297	0.284	0.296
$W \sim f_{-i}(W)$		0.370	0.215	0.163	0.414	0.232	0.340	0.242	0.335	0.319	0.264
Panel III. Own Xs, own Ws, $\varepsilon=0$, and											
$\eta \sim f(\eta)$	0.444	0.430	0.452	0.438	0.448	0.447	0.435	0.437	0.446	0.438	0.447
$\eta \sim f_i(\eta)$		0.430	0.451	0.439	0.447	0.447	0.436	0.449	0.441	0.447	0.442
$\eta \sim f_{-i}(\eta)$		0.431	0.450	0.437	0.449	0.445	0.436	0.433	0.457	0.433	0.456
Panel IV. Own α, $\varepsilon=0$, and											
$X=E[X]$	0.607	0.565	0.632	0.622	0.594	0.623	0.568	0.630	0.597	0.591	0.616
$X \sim f(X)$	0.597	0.564	0.616	0.607	0.588	0.612	0.559	0.622	0.586	0.589	0.601
$X \sim f_i(X)$		0.581	0.605	0.605	0.591	0.606	0.575	0.595	0.598	0.597	0.596
$X \sim f_{-i}(X)$		0.554	0.633	0.608	0.587	0.630	0.551	0.634	0.556	0.584	0.610
Panel V. Own Xs, own α, and											
$\varepsilon \sim f(\varepsilon)$	0.597	0.572	0.612	0.619	0.580	0.605	0.577	0.586	0.601	0.598	0.596
$\varepsilon \sim f_i(\varepsilon)$		0.572	0.612	0.611	0.588	0.606	0.576	0.589	0.600	0.599	0.597
$\varepsilon \sim f_{-i}(\varepsilon)$		0.574	0.611	0.628	0.571	0.603	0.580	0.585	0.605	0.597	0.597
Panel VI. Own α and											
$X, \varepsilon \sim f(X, \varepsilon)$	0.589	0.559	0.607	0.599	0.581	0.603	0.553	0.614	0.578	0.581	0.593
$X, \varepsilon \sim f_i(X, \varepsilon)$		0.573	0.596	0.591	0.590	0.597	0.569	0.590	0.590	0.589	0.588
$X, \varepsilon \sim f_{-i}(X, \varepsilon)$		0.548	0.622	0.606	0.573	0.620	0.546	0.625	0.554	0.575	0.602

Notes: Benchmark case denotes the observed probability in the sample. Simulations - based on 500 draws - obtained after estimating the dynamic model $y_{it} = \alpha_i + \gamma y_{it-1} + X_{it}\beta + \varepsilon_{it}$, where $\alpha_i = w_i\delta + \eta_i$. $f(\cdot)$ denotes the empirical distribution of the argument. Sample includes data from waves 1-4 of the ECLS-B. See text for the list of covariates and further details.

Table 19. Dynamic Simulations: Height, $\Pr(y_{it} \geq 95^{\text{th}} \text{ percentile} \mid y_{it} \geq 95^{\text{th}} \text{ percentile})$.

	Full	Race		Gender		Urban Status		Education		SES	
	Sample	White	Non-White	Male	Female	Urban	Non-Urban	Non-College	College	Low SES	High SES
Benchmark	0.365	0.344	0.377	0.372	0.360	0.355	0.387	0.353	0.370	0.319	0.393
Panel I. Own Xs, $\varepsilon=0$, and											
$\alpha=E[\alpha]$	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
$\alpha \sim f(\alpha)$	0.092	0.102	0.087	0.061	0.112	0.088	0.100	0.080	0.096	0.095	0.090
$\alpha \sim f_i(\alpha)$		0.080	0.100	0.074	0.094	0.094	0.082	0.085	0.093	0.090	0.091
$\alpha \sim f_{-i}(\alpha)$		0.116	0.069	0.050	0.128	0.072	0.105	0.077	0.101	0.095	0.086
Panel II. Own Xs, $\eta=0$, $\varepsilon=0$, and											
$W=W_i$	0.020	0.008	0.027	0.020	0.020	0.021	0.017	0.015	0.022	0.022	0.019
$W \sim f(W)$	0.007	0.007	0.007	0.002	0.010	0.006	0.008	0.004	0.008	0.007	0.007
$W \sim f_i(W)$		0.004	0.008	0.003	0.005	0.007	0.006	0.005	0.007	0.007	0.006
$W \sim f_{-i}(W)$		0.009	0.004	0.001	0.015	0.005	0.008	0.004	0.009	0.008	0.006
Panel III. Own Xs, own Ws, $\varepsilon=0$, and											
$\eta \sim f(\eta)$	0.136	0.131	0.139	0.125	0.144	0.134	0.141	0.129	0.139	0.133	0.138
$\eta \sim f_i(\eta)$		0.125	0.141	0.127	0.138	0.132	0.139	0.125	0.137	0.137	0.133
$\eta \sim f_{-i}(\eta)$		0.134	0.131	0.120	0.146	0.129	0.142	0.129	0.137	0.128	0.142
Panel IV. Own α, $\varepsilon=0$, and											
$X=E[X]$	0.350	0.320	0.366	0.346	0.354	0.369	0.309	0.426	0.322	0.297	0.382
$X \sim f(X)$	0.353	0.332	0.365	0.360	0.349	0.368	0.322	0.416	0.330	0.298	0.386
$X \sim f_i(X)$		0.346	0.356	0.358	0.353	0.362	0.337	0.388	0.341	0.305	0.383
$X \sim f_{-i}(X)$		0.322	0.380	0.362	0.348	0.384	0.315	0.429	0.304	0.297	0.395
Panel V. Own Xs, own α, and											
$\varepsilon \sim f(\varepsilon)$	0.304	0.306	0.303	0.311	0.300	0.301	0.311	0.308	0.303	0.264	0.328
$\varepsilon \sim f_i(\varepsilon)$		0.303	0.304	0.302	0.307	0.302	0.310	0.308	0.302	0.267	0.326
$\varepsilon \sim f_{-i}(\varepsilon)$		0.307	0.301	0.319	0.292	0.298	0.310	0.306	0.304	0.263	0.330
Panel VI. Own α and											
$X, \varepsilon \sim f(X, \varepsilon)$	0.357	0.336	0.368	0.364	0.353	0.372	0.325	0.415	0.335	0.306	0.388
$X, \varepsilon \sim f_i(X, \varepsilon)$		0.349	0.361	0.358	0.360	0.365	0.339	0.392	0.345	0.315	0.382
$X, \varepsilon \sim f_{-i}(X, \varepsilon)$		0.328	0.381	0.373	0.346	0.385	0.320	0.425	0.313	0.301	0.398

Notes: See Table 18 and text for further details.

Table 20. Dynamic Simulations: Height, $\Pr(y_{it} \geq 85^{\text{th}} \text{ percentile} \mid y_{it} \leq 50^{\text{th}} \text{ percentile})$.

	Full	Race		Gender		Urban Status		Education		SES	
	Sample	White	Non-White	Male	Female	Urban	Non-Urban	Non-College	College	Low SES	High SES
Benchmark	0.179	0.176	0.181	0.168	0.193	0.185	0.165	0.200	0.170	0.196	0.169
Panel I. Own Xs, $\varepsilon=0$, and											
$\alpha=E[\alpha]$	0.183	0.206	0.163	0.071	0.319	0.160	0.239	0.117	0.210	0.218	0.162
$\alpha \sim f(\alpha)$	0.383	0.401	0.368	0.345	0.428	0.373	0.407	0.357	0.393	0.393	0.377
$\alpha \sim f_i(\alpha)$		0.350	0.401	0.381	0.389	0.385	0.378	0.380	0.383	0.387	0.380
$\alpha \sim f_{-i}(\alpha)$		0.436	0.319	0.310	0.466	0.342	0.419	0.348	0.418	0.396	0.370
Panel II. Own Xs, $\eta=0$, $\varepsilon=0$, and											
$W=W_i$	0.148	0.111	0.178	0.139	0.157	0.156	0.127	0.125	0.157	0.145	0.149
$W \sim f(W)$	0.296	0.321	0.274	0.231	0.375	0.280	0.334	0.255	0.313	0.314	0.285
$W \sim f_i(W)$		0.254	0.315	0.280	0.316	0.300	0.278	0.278	0.302	0.291	0.296
$W \sim f_{-i}(W)$		0.369	0.215	0.180	0.431	0.231	0.357	0.245	0.340	0.326	0.264
Panel III. Own Xs, own Ws, $\varepsilon=0$, and											
$\eta \sim f(\eta)$	0.297	0.276	0.314	0.292	0.303	0.301	0.288	0.283	0.303	0.293	0.299
$\eta \sim f_i(\eta)$		0.273	0.318	0.295	0.302	0.303	0.287	0.288	0.301	0.302	0.295
$\eta \sim f_{-i}(\eta)$		0.281	0.310	0.288	0.307	0.300	0.288	0.282	0.308	0.288	0.309
Panel IV. Own α, $\varepsilon=0$, and											
$X=E[X]$	0.176	0.158	0.191	0.179	0.172	0.189	0.144	0.200	0.166	0.168	0.181
$X \sim f(X)$	0.204	0.186	0.219	0.207	0.201	0.208	0.193	0.230	0.193	0.202	0.205
$X \sim f_i(X)$		0.194	0.212	0.205	0.203	0.204	0.205	0.208	0.201	0.207	0.201
$X \sim f_{-i}(X)$		0.179	0.229	0.207	0.200	0.219	0.187	0.239	0.174	0.199	0.210
Panel V. Own Xs, own α, and											
$\varepsilon \sim f(\varepsilon)$	0.158	0.148	0.166	0.159	0.156	0.160	0.154	0.162	0.156	0.170	0.150
$\varepsilon \sim f_i(\varepsilon)$		0.146	0.167	0.154	0.161	0.160	0.150	0.163	0.156	0.173	0.149
$\varepsilon \sim f_{-i}(\varepsilon)$		0.150	0.163	0.164	0.151	0.158	0.154	0.163	0.155	0.168	0.153
Panel VI. Own α and											
$X, \varepsilon \sim f(X, \varepsilon)$	0.214	0.197	0.229	0.217	0.212	0.218	0.204	0.241	0.203	0.213	0.215
$X, \varepsilon \sim f_i(X, \varepsilon)$		0.204	0.224	0.210	0.219	0.215	0.214	0.220	0.211	0.220	0.211
$X, \varepsilon \sim f_{-i}(X, \varepsilon)$		0.193	0.237	0.223	0.205	0.228	0.201	0.249	0.186	0.210	0.223

Notes: See Table 18 and text for further details.